

UNCLASSIFIED

Hanford

HW-64371

AEC RESEARCH AND DEVELOPMENT REPORT

EVALUATION OF RADIOLOGICAL CONDITIONS IN THE VICINITY OF HANFORD FOR 1959

R. L. JUNKINS
E. C. WATSON
I. C. NELSON
R. C. HENLE

MAY 9, 1960

HANFORD LABORATORIES

HANFORD ATOMIC PRODUCTS OPERATION
RICHLAND, WASHINGTON

GENERAL  ELECTRIC

UNCLASSIFIED

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

UNCLASSIFIED

HW-64371

UC-41, Health and Safety
(TID-4500, 15th Ed.)

EVALUATION OF RADIOLOGICAL CONDITIONS
IN THE VICINITY OF HANFORD FOR 1959

By

The Radiological Evaluation Staff

R. L. Junkins

E. C. Watson, I. C. Nelson and R. C. Henle
Radiation Protection Operation

May 9, 1960

HANFORD ATOMIC PRODUCTS OPERATION
RICHLAND, WASHINGTON

Work performed under Contract No. AT(45-1) - 1350 between
the Atomic Energy Commission and General Electric Company

Printed by/for the U. S. Atomic Energy Commission

Printed in USA. Price \$2.50. Available from the
Office of Technical Services
U. S. Department of Commerce
Washington 25, D. C.

UNCLASSIFIED

TABLE OF CONTENTS

	<u>Page</u>
I. INTRODUCTION	4
II. SUMMARY	5
III. DISCUSSION OF ENVIRONMENTAL SAMPLING RESULTS	6
A. Concentrations of Radionuclides in the Columbia River	6
Sanitary Water Concentrations	6
Concentrations in Edible Fish and Waterfowl.	19
Recreational Use of Columbia River	22
Concentrations in Columbia River Water at Vancouver.	22
Results Related to the Pacific Ocean	23
B. Concentration of Radionuclides in Gaseous Effluents and	
Vegetation	26
Atmospheric Concentrations.	26
Concentrations in Vegetation	32
C. Concentration of Radionuclides in Farm Products	41
Concentrations in Milk	44
Concentrations in Fresh Farm Produce	47
Radionuclides in Miscellaneous Foodstuffs	52
IV. ESTIMATE OF RADIATION EXPOSURE - NONOCCUPATIONAL .	53
V. ACKNOWLEDGEMENTS	56
BIBLIOGRAPHY	57
APPENDIX A - ANALYTICAL METHODS	59
APPENDIX B - RIVER AND RELATED SAMPLE RESULTS	65
APPENDIX C - ATMOSPHERIC AND VEGETATION SAMPLE	
RESULTS	82
APPENDIX D - FARM PRODUCT SAMPLE RESULTS	108

LIST OF FIGURES

<u>No.</u>	<u>Title</u>	<u>Page</u>
1	Hanford Plant Environs	7
2	Isotopic Composition of Reactor Effluent 4 Hours and 24 Hours after Irradiation	8
3	Reduction in Radioactivity of Reactor Effluent with Time	9
4	Depletion of Several Radionuclides in Columbia River Water During Treatment at Pasco, Washington Water Plant (1959 Averages)	11
5	Depletion of Selected Radionuclides in Columbia River Water During Treatment at Pasco, Washington Water Plant (1959 Monthly Averages)	12
6	Per Cent of MPC_w for the GI Tract for Selected Radionuclides in Columbia River Water and Sanitary Water at Pasco, Washington - 1959	14
7	Variations in Concentrations of Selected Radionuclides in Columbia River Water at Pasco, Washington for 1959	15
8	Flow Rate of Columbia River at Pasco, Washington	16
9	Volumes of Water and Sewage Processed at Pasco, Washington Water and Sewage Plants 1959 Monthly Averages	17
10	Concentrations of Beta Emitters in Muscle of Whitefish at Ringold	21
11	Columbia River Flow at Pasco, and Vancouver, Washington-1959	24
12	Radioactive Particle Concentrations at Selected Northwestern U. S. Sampling Locations - 1959	27
13	Activity on Filters from Selected Northwestern U. S. Sampling Locations - 1959	28
14	Comparison of Particle Distribution Found on Filter Paper- 4th Quarter 1959	30
15	Vegetation Sampling Zone Locations-Hanford and Environs	33
16	Radionuclide Concentration on Native Grasses-Zone C (1959)	34
17	Radionuclide Concentration on Native Grasses-Zone O (1959)	35
18	Radionuclide Concentration on Native Grasses-Spokane and Vicinity (1959)	36
19	Radionuclide Concentration on Native Grasses-Portland and Vicinity (1959)	37
20	Zr ⁹⁵ -Nb ⁹⁵ Concentration on Native Grasses at Selected Sites-1959	39
21	Comparison of I ¹³¹ Deposition on Grasses with I ¹³¹ Emitted in the Vicinity of a Separations Facility - 1959	40
22	Comparison of Average I ¹³¹ Concentrations - Jack Rabbit Thyroids and Native Grasses (1959)	42

EVALUATION OF RADIOLOGICAL CONDITIONS
IN THE VICINITY OF HANFORD FOR 1959

I. INTRODUCTION

During operation of the plutonium production and research facilities at Hanford, * controlled amounts of radioactive wastes are released to the atmosphere, ground, and to the Columbia River. These wastes contribute to the radiation exposure of persons living in the neighborhood of the controlled area. The protection of these persons from undue radiation exposure attributable to Hanford sources is one of the attendant responsibilities in the operation of the Hanford facilities.

The recommendations of the NCRP and ICRP, and the results of a continuous research program,⁽¹⁾ form the basis of the standards used in assessing the degree of protection needed and attained. The effectiveness of waste control and radiation protection practices is determined by comparison of the results of an extensive program of sampling and measurement of radionuclide abundance in the Hanford environs with the reference standards.

The release of radioactive wastes and environmental radiation protection results through 1958 are summarized in the transcribed proceedings of the Joint Committee on Atomic Energy - Congressional Hearings held in 1959.⁽²⁾

Hanford's experience in the field of environmental radiation protection for the year 1959 is the subject of this report. The results of Hanford's environmental monitoring for 1958 are reported in "Hanford Environmental Monitoring Annual Report", HW-61676, Unclassified, August 27, 1959, by B. V. Andersen.

* Operated for the Atomic Energy Commission by the General Electric Company under contract number AT(45-1)-1350.

II. SUMMARY

A large fraction of exposure to persons residing in the neighborhood of the Hanford controlled area is through drinking treated Columbia River water. The source of this exposure is identified with the neutron-induced radionuclides generated in the cooling water of the production reactors. However, other sources and paths of exposure contribute to the total dose estimate.

The environmental radiation exposure is estimated for each of the critical organs of major interest, i. e., the gastrointestinal tract, bone and the total body. In each case, multiple sources and paths of intake contribute in varying degrees to the estimated dose. There is a general lack of detailed information on the variation of diet within a population group. Because of these and other uncertainties the environmental radiation exposure cannot be stated precisely. However, the exposure to persons in the neighborhood of the Hanford controlled area is estimated to be about 10 per cent-15 per cent of that permitted by the recommendations of the NCRP.

The estimated dose consists primarily of the dose due to ingestion of drinking water and foodstuffs which have been irrigated by river water containing diluted reactor cooling water. Hanford's contribution to environmental exposure through atmospheric paths is considerably less than that due to fallout from nuclear detonations. Atmospheric contributions are small compared to the contribution through water effluents.

It is possible that for exceptional cases where unusual amounts of local fish are included in the diet, the exposure would be greater than the preceding estimate. This larger estimate appears to be contained within the range of 40 per cent-60 per cent of the appropriate maximum permissible limit.

III. DISCUSSION OF ENVIRONMENTAL SAMPLING RESULTS

A brief discussion of the analytical methods for determining radionuclide concentrations in the various types of samples is contained in Appendix A. The results and interpretation thereof as discussed in this section are categorized according to method of waste release and potential path of intake.

A. Concentrations of Radionuclides in the Columbia River

The eight production reactors located as shown in Figure 1 create radionuclides in that portion of the Columbia River water used as reactor coolant. More than 60 radionuclides have been measured in reactor effluent water, and their relative abundance is illustrated in Figure 2. Other Hanford facilities do not contribute measurably to the quantity of radionuclides in the river.

Many of the radionuclides in the reactor effluent water are short-lived and decay rapidly after formation in the reactors. A gross decay curve, which illustrates this property, is reproduced as Figure 3. In addition to radioactive decay, a quantity of the radionuclides is removed from the river water by such mechanisms as silting and uptake by river biota. (3)

There are several ways by which radionuclides in the river water may result in exposure to humans. Among these paths of exposure are: 1) irrigated agriculture products (discussed in Section C), 2) sanitary water derived from treating river water, 3) edible fish and waterfowl which inhabit the river, and 4) external exposure from use of the river for recreational purposes.

Sanitary Water Concentrations

Pasco and Kennewick are the nearest cities downstream from the Plant which treat Columbia River water for use as sanitary water. Sanitary water from each of the water treatment plants was sampled weekly and analyzed for individual radionuclides or in some cases groups of radionuclides





FIGURE 1



In addition to the isotopes shown above, which contribute about 98 per cent of the activity 4 hours after irradiation, trace amounts of the following have also been found.

Eu-152	Eu-157	I-131*	Y-91	Pr-145	Cs-137
Sm-153	Ba-140*	Ce-141	Fe-59*	Pm-151	Sr-85
W-187	Mo-99	Pr-142	Sr-89*	Co-60	U-238*
La-141	Sm-156	C-14	Mn-54	Pr-143	Pu-239*
Nd-149	Sc-46*	Nd-147	Zr-95	Ru-103	Ac-227
La-140	Cd-115	Ca-45*	Pm-149	Sc-47	Po-210*
I-132*	Ce-143	Ag-111	Eu-156	Sr-90*	

* Routine measurements are made on these isotopes.

FIGURE 2
Isotopic Composition of Reactor Effluent
4 Hours and 24 Hours after Irradiation

REDUCTION IN RADIODACTIVITY OF REACTOR EFFLUENT WITH TIME




FIGURE 3

such as the rare earths. Untreated river water at the Pasco pumping plant inlet was also sampled and analyzed weekly. The data are included in Appendices B-1, -2, and -3.

The sanitary water of other communities near the plant which have sources of water supply other than the Columbia River was periodically sampled and analyzed. Richland, Benton City, Ringold and Riverview farms are examples of this portion of the program. A higher concentration of natural uranium has been noted in the water from Benton City wells, than in the water from wells of other local communities throughout a ten year sampling period. See Appendices B-4 and -5. However, there is no evidence that this condition is attributable to Hanford facilities. Sampling and analyses of the river and sanitary water in the Pasco system and sanitary water in Kennewick provided data which were used to estimate the radiation exposure from this source to the local residents. In both cities, the sanitary water samples were collected near the water treatment plants. Because there is an unknown, but significant flow time between sampling locations and most consumers, the concentration in sanitary water at the time it is consumed is overestimated. This is due to radioactive decay which reduces the concentrations of some of the nuclides substantially before they reach the majority of the consumers.

The actual decay time available varies from hours to days depending upon water usage rates, particular location of the consumer, and other influences.⁽⁴⁾ Figures 4 and 5 show the efficiency of the water treatment plant at Pasco for the removal of various radionuclides. These data include the radioactive decay of the radionuclides during travel through the water treatment plant.

For the mixture of nuclides present in the river water and sanitary water, the gastrointestinal (GI) tract is an organ of major interest. During 1959, the calculated average dose to the GI tract for the Pasco population was equivalent to ~ 5 per cent of the maximum permissible concentration in water (MPC_w) for persons in the neighborhood of controlled areas.⁽⁵⁾ The average monthly results for 1959 are summarized in Table I.




FIGURE 4

Depletion of Several Radionuclides in Columbia River Water During Treatment at Pasco, Washington Water Plant (1959 Averages)




FIGURE 5

Depletion of Selected Radionuclides in Columbia River Water During Treatment
at Pasco, Washington Water Plant (1959 Monthly Averages)

TABLE I
AVERAGE PERCENTAGES OF NBS HANDBOOK 69 MPC_W -
GI TRACT IN SANITARY WATER AT PASCO, WASHINGTON 1959

Month	% MPC _W *
January	5.8
February	5.3
March	5.4
April	5.2
May	3.7
June	2.1
July	2.0
August	4.3
September	4.0
October	6.2
November	3.3
December	4.5

* The MPC_W for the GI tract was taken as that given for continuous occupational exposure from NBS Handbook 69 and multiplied by 0.1 for persons residing in the neighborhood of a controlled area.
Solubility in GI tract and body fluids is assumed.

The calculated per cent for Kennewick residents averaged ~ 2 per cent of the MPC_W for the GI tract. Although there was no known routine consumption of the untreated river water, such a source of drinking water throughout the year would have resulted in radiation exposure equivalent to ~ 15 per cent of the MPC_W for the GI tract. The average contribution of several radionuclides to the dose to the GI tract would be as illustrated in Figure 6. Figure 7 illustrates the seasonal variation in concentration of radionuclides. This variation is influenced by the river flow variation shown in Figure 8, and, in the case of sanitary water, the treatment plant throughput ⁽⁴⁾ illustrated in Figure 9.




FIGURE 6

Per Cent of MPC_W for the GI Tract for Selected Radionuclides in Columbia River
Water and Sanitary Water at Pasco, Washington - 1959




FIGURE 7
Variations in Concentrations of Selected Radionuclides in Columbia River Water
at Pasco, Washington for 1959




FIGURE 8

Flow Rate of Columbia River at Pasco, Washington
(From Data Published by the U. S. Geological Survey)




FIGURE 9

Volumes of Water and Sewage Processed at Pasco, Washington
Water and Sewage Plants
1959 Monthly Averages

In calculating the percentage MPC_w for the GI tract from drinking water, the rare earth elements are included as a group; a maximum permissible limit for which was derived from measured concentrations in reactor effluent water⁽⁶⁾ and the individual MPC_w 's.⁽⁵⁾

The estimated dose to other organs through water intake is substantially less than that for the GI tract. Some indicative values are compared with the GI tract exposure in Table II.

TABLE II
1959 AVERAGE PERCENTAGE MPC_w^* FOR ORGANS OF INTEREST

	% MPC_w		
	<u>Bone</u>	<u>Total Body</u>	<u>GI</u>
Pasco, Inlet Water (Columbia River Water)	1.6	0.9	15
Pasco, Sanitary Water	0.9	0.5	5
Kennewick, Sanitary Water	0.6	0.2	2
Vancouver, Columbia River Water**	0.7	0.2	-

* NBS Handbook 69, See footnote Table I.

** Last six months of 1959.

Several radionuclides which contribute to the exposure to these organs are present in amounts below the analytical detection limits. In this case and where analytical data permit, the concentrations in the reactor effluent, river depletion rates and river flow data are used to estimate the concentration at a given location. If insufficient data are available for this computation, it is then assumed that the radionuclide was present in the concentration represented by the detection limit. Thus the values in Table II are believed to be upper bounds of the estimated exposure.

The percentage of MPC_w for other organs was less than those listed in the preceding table.

The filter backwash solids and sediments accumulated in settling basins in Pasco water plant were sampled intermittently throughout the year and analyzed for gross beta emitters. These solids are dispersed into the Columbia

River along with waste water. Measurements⁽⁴⁾ made during 1959 indicate this path of exposure to downstream communities to be negligible.

A study was made of distribution of radionuclides in the Pasco sewage plant, particularly the digested solids. In all of these samples the long-lived radionuclides predominated; Zn⁶⁵ and Cr⁵¹ were the most abundant. Shorter-lived radionuclides decayed to below detectable concentrations during the process period. In the dried solids from the digested sewage, the Zn⁶⁵ concentration was $\sim 5 \times 10^{-5} \mu\text{c}$ Zn⁶⁵/g. Chromium-51 was the second most abundant nuclide but was an order of magnitude less in concentration.

It is of interest to estimate the uptake of Zn⁶⁵ and Cr⁵¹ by plants should such plants be fertilized by these solids. This estimate assumes: 1) the concentration remains the same from the time of production to the time of actual use, 2) that 10⁻⁵ grams of this fertilizer are used per gram of soil available to the plant (about twice average application)⁽⁷⁾, and 3) rates of uptake equivalent to 1 μc Zn⁶⁵/g dry soil results in 0.01 μc Zn⁶⁵/g dry leaf and 1 μc Cr⁵¹/g dry soil yields 0.2 μc Cr⁵¹/g dry leaf.⁽⁷⁾ The estimated concentrations in leaves of vegetation is: $\sim 5 \times 10^{-12} \mu\text{c}$ Zn⁶⁵/g dry leaf and $\sim 10^{-11} \mu\text{c}$ Cr⁵¹/g dry leaf. Both of these concentrations are sufficiently small to make detection difficult and they are negligible in terms of exposure.

Concentrations in Edible Fish and Waterfowl

The Ringold vicinity is the first downstream area accessible to the public. It is a sportsfishing area where whitefish are taken, particularly during the fall and winter months. Waterfowl inhabit this section of the river; and some of these are harvested by local hunters. The amount of radiation exposure received by persons who regularly harvest ducks and whitefish was recently estimated by Foster and Junkins.⁽⁸⁾ The following excerpt from the referenced document describes the present situation:

"The whitefish is currently considered to be the species which may contribute most to human exposure because of its relatively high accumulation of radionuclides (only slightly less than suckers), and because it is easily caught during the fall at a time when its content of radioactive materials is maximum.

"The maximum permissible concentration for P³² in drinking water 'for persons in the neighborhood of controlled areas' derived from NBS Handbook 69 is $2 \times 10^{-5} \mu\text{c}/\text{cc}$ which is equivalent to a daily intake of $4.4 \times 10^{-2} \mu\text{c}/\text{day}$. This amounts to about 0.3 μc per week or about 16 μc per year.

"On the basis of the stated assumptions and with averaging over a year's period, this quantity of P³² could be obtained from the consumption of about one pound of whitefish flesh each week in years when the average concentration of P³² in the fish reaches about $2 \times 10^{-3} \mu\text{c}/\text{g}$ of flesh during the late summer or early fall months.

"The concentration of Zn⁶⁵ in the whitefish during the last summer is only about 1/20 of the observed level for P³² (in August, 1957). Further, the maximum permissible intake rate for Zn⁶⁵ is 5-fold greater than for P³². The Zn⁶⁵ thus contributes only a relatively small fraction of the exposure received from the consumption of fish.

"The P³² content of the flesh of some ducks which are killed along the river within the Hanford reservation is similar to that found in the fish. A comparatively few ducks remain on the river throughout the year so that the accumulation of P³² and a few other nuclides builds up to a level which is in equilibrium with their environment. These ducks are not available to off-project hunters until the fall hunting season at which time their number is diluted 200 to 1,000 fold by flocks of ducks which are migrating through the region."

Results of the 1959 biological monitoring program, conducted by Radioecology Operation, indicated lower concentrations of nuclides in fish and waterfowl than were observed in 1957 and 1958. (9, 10, 11) The data are contained in Appendix B-6. Measurements made during the past four years on fish sampled at Ringold are summarized in Figure 10.




FIGURE 10
Concentrations of Beta Emitters
in Muscle of Whitefish at Ringold

Recreational Use of Columbia River

In the vicinity of Richland, Pasco and Kennewick, ionization chambers were used to measure the penetrating radiation in the river. Such measurements indicated an annual exposure of about 60 mr, excluding background, for the operators of a ferry boat just upstream from the City of Richland, assuming they spent one-third of their time in the boat. Since the major contributors to the dose, such as Na^{24} , are short-lived, the exposure from recreational boating further downstream is less. Swimmers would have received about 6 mr from this source, if they were in the river as much as 240 hours during the year.

Concentrations in Columbia River Water at Vancouver

Most of the radionuclides present in the river water which contribute to the GI tract dose have decayed to concentrations too small to be measured at Vancouver, Washington. Radionuclides which contribute to the dose to bone and total body have decayed to a lesser extent and the calculated percentage of the respective MPC_w 's in river water at Vancouver were all of the same order of magnitude; i. e., < 1 per cent of the applicable MPC_w . The measurements at Vancouver are reported in Appendix B-7 and are summarized in Table III.

TABLE III

1959 AVERAGE PERCENTAGE MPC_w * FOR SELECTED ORGANS FROM DRINKING COLUMBIA RIVER WATER AT VANCOUVER, WASHINGTON

	% MPC_w		
	GI	Bone	Total Body
P^{32}	0.034	0.15	0.034
Cr^{51}	0.11	--	0.011
Zn^{65}	0.014	0.0029	0.029
Np^{239}	0.018	~0	~0

* NBS Handbook 69 (See * Table I)

There is considerable uncertainty in the related estimates because of the low concentrations of the radionuclides; however, the results are consistent with those calculated from upstream measurements of higher concentrations.

Samples of river water and mud were collected in several locations downstream from Pasco and are reported in Appendices B-8, -9 and 10. These samples were analyzed for gross alpha and beta emitters and the data used as an indicator of unusual conditions. During 1959, analyses for individual radionuclides were initiated on the samples collected at Vancouver.

Results Related to the Pacific Ocean

The analytical measurements together with information on the river flow, Figure 11, can be used to estimate the total quantities of the various radionuclides transported to the Pacific Ocean by the river. Such estimates are included in Appendices B-11 and -12. Chromium-51, Np^{239} , Zn^{65} , and P^{32} persisted in measurable amounts at Vancouver. Although the half-lives of some of these radionuclides are long by comparison with most of the others in the mixture, they are not so long that the total quantities in the North Pacific Ocean will continue to increase indefinitely. On the basis of the Vancouver measurements, the total quantity of these radionuclides in the Ocean, attributable to the Hanford reactors may be estimated. If a constant rate of entry into the ocean, over a period of years, equivalent to that indicated by the Vancouver data is assumed, then the equilibrium value amounts to: ~400 curies of P^{32} , ~400 curies of Np^{239} , ~40,000 curies of Cr^{51} and ~7,000 curies of Zn^{65} . An equilibrium exists for these radionuclides where the rate of addition through the river system corresponds to the rate of decay of the radionuclides which have previously entered the Ocean.

Vancouver, Washington was selected as the farthest downriver location for routine sampling for several reasons: The salt content of

COLUMBIA RIVER FLOW AT PASCO AND VANCOUVER - 1959
FROM DATA PUBLISHED BY THE U. S. GEOLOGICAL SURVEY




FIGURE 11

water nearer the mouth of the river complicates quantitative measurement of the radionuclides; tidal movement near the mouth increases variability of results and interpretation of flow times; and, since the very short-lived radionuclides have decayed to small concentrations, radioactive decay of the remainder decreases the total radioactivity at much slower rate than is typical for upriver locations.

The most probable way in which these radionuclides can contribute to human exposure is through concentration in edible marine organisms taken from the Ocean.

Concentrations of the radionuclides in fish, shell fish, and other biota were measured in samples collected along the coasts of Oregon and Washington.

The measurement of Zn⁶⁵ in people has been reported previously. (13) Since Zn⁶⁵ was known to exist in some foodstuffs grown in the Riverview District, it was suspected that people whose diets included these foods would also contain measurable amounts of Zn⁶⁵. This was confirmed in 1958. (14) However, in the course of calibrating and refining Hanford's whole body counter facility, it became apparent that there was yet another source of Zn⁶⁵ intake. (15) This source has since been identified as sea food, primarily oysters, grown in the Pacific Ocean.

Results of sea food sampling and analyses are contained in Appendix B-13, -14 and -15. The data indicate that the oysters, particularly those from the vicinity of Willapa Bay, Washington, were higher in Zn⁶⁵ content than were other sea foods. These oysters contained on the order of 3.0 to $7.0 \times 10^{-5} \mu\text{c}$ Zn⁶⁵ ~g of edible meat. Measured body burdens of those whose diets are known to include these oysters were 0.01 μc Zn⁶⁵ or less, which is < 0.2 per cent of the maximum permissible body burden.

Since the Columbia River transports Zn⁶⁵ to the Ocean, it might be expected that Zn⁶⁵ concentrations in marine organisms would vary along the coast, relative to the mouth of the river. Due to the unavailability of some organisms at various locations, it is difficult to establish such a pattern.

However, it is interesting to note that the Zn⁶⁵ content of oysters from Sequim, Washington, which is on the Straits of Juan de Fuca, is about one order of magnitude less than those of Willapa Bay, Washington.

B. Concentration of Radionuclides in Gaseous Effluents and Vegetation

Any discussion of concentrations of radionuclides in the atmosphere or on vegetation during 1959 must include consideration of the contribution of fallout of debris from nuclear detonations. In the United States, the testing of nuclear devices by detonation was terminated for an indefinite period during the latter part of 1958. However, the fallout of debris from prior detonations continued to be measurable throughout the early months of 1959. During these early months of 1959, the contribution from fallout was sufficient to make the detection of any contribution normally associated with operation of Hanford's facilities difficult.

Atmospheric Concentrations

The filter papers from air sampling stations were changed daily by cooperating agencies. These filter papers were analyzed both for number of radioactive particles and for total beta activity. The average weekly particle concentrations measured in number of particles per cubic meter of filtered air are contained in Appendix C-1 and illustrated in Figure 12. The results for total beta emitters are reported in Appendix C-2 and illustrated in Figure 13. The trend throughout the year in both Figures 12 and 13 is apparent. It is indicated by a decrease in the average weekly concentration and the average weekly beta activity beginning in April. The decrease is logarithmic over the next four to five months, and reaches an apparent "background" in October.

During the period from October through December, 1959 measurements of filter papers indicated a measurable contribution from Hanford Operations.




FIGURE 12
Radioactive Particle Concentrations at Selected Northwestern United States
Sampling Locations - 1959




FIGURE 13

Activity on Filters from Selected Northwestern United States Sampling Locations - 1959

This contribution was primarily limited to the Hanford controlled area as demonstrated by the plots of the particle frequency distribution. Only those filters through which equal volumes of air (i.e., $\sim 120 \text{ m}^3$) have been passed are considered. Figure 14b shows such a distribution for the location within the controlled area identified as "Hanford" and Figure 14a is a plot for all off-site locations. The distribution for "Hanford" is very nearly described by a Poisson distribution with a mean of about one particle per 24-hour sampling period. Air sample filters for locations distant from Hanford had a particle distribution, which again could be described by a Poisson distribution but with a mean of 0.4 particles per 24-hour sampling period. The discrepancy, as indicated by marginal χ^2 test results can be explained by noting that the observer, in counting the radiographic images, either tends to find no images (particles) or, once he has found one, he tends to find more. Thus, the distribution shows a tendency to be high at zero as well as at the larger numbers of particles counted; such distributions have been described by a negative binomial distribution. ⁽¹⁶⁾ This effect has been observed by other investigators in dealing with the counting of a very small number of events per unit area (or time) by human observers.

The analysis of filter papers for individual radionuclides is not a routine procedure. However, such an analysis was performed ⁽¹⁷⁾ in 1959. During the third calendar quarter of 1959, filter papers from each of 13 locations were grouped for this analysis. Table IV is a summary of the results corrected for radioactive decay to July 15, 1959 and converted to μc per cc of filtered air. The fact that the concentrations in Table IV are very nearly the same for all locations is further evidence that the radionuclides are from fallout.




FIGURE 14
Comparison of Particle Distribution Found on Filter Paper
4th Quarter - 1959

TABLE IV
CONCENTRATION OF RADIONUCLIDES ON FILTER PAPERS
THIRD QUARTER - 1959 (Units of $10^{-13} \mu\text{c}/\text{cc}$)

	<u>Zr</u> ⁹⁵ - <u>Nb</u> ⁹⁵	<u>Cs</u> ¹³⁷	<u>Ru</u> ¹⁰⁶	<u>Ce</u> ¹⁴⁴ - <u>Pr</u> ¹⁴⁴	<u>Zn</u> ⁶⁵
Benton City, Wash.*	1.4	0.47	1.6	2.9	--
Richland, Wash.	1.3	0.29	1.7	2.5	--
Kennewick, Wash.*	1.9	0.54	2.1	3.8	--
Pasco, Wash.*	1.7	0.53	2.3	3.4	0.11
Yakima, Wash.	1.7	0.27	1.5	2.7	--
Walla Walla, Wash.	1.4	0.29	1.5	2.9	--
Meacham, Ore.*	0.97	0.31	1.1	2.6	0.09
Lewiston, Idaho	1.6	0.35	2.7	3.7	--
Spokane, Wash.	1.4	0.29	1.6	3.1	--
Boise, Idaho	2.0	0.44	2.7	5.0	--
Seattle, Wash.*	0.90	0.25	0.88	1.8	0.06
Klamath Falls, Ore.	1.4	0.32	2.2	3.0	--
Great Falls, Mont.*	1.4	0.37	1.5	2.5	0.11
Average Concentration	<u>1.47</u>	<u>0.363</u>	<u>1.80</u>	<u>3.07</u>	<u>0.09</u>
Per Cent MPC _{air} for Critical Organ **	0.001	0.002	0.004	0.01	(0.0002)***

* Not routinely analyzed for particle concentration.

** Based on NCRP recommendations.

*** Based on four locations with reported results.

Radiiodine has characteristically predominated in gaseous effluents from Hanford separations facilities. Therefore, air sampling for I¹³¹ in the atmosphere is a routine procedure for several nearby communities. The sampling results are given in Appendix C-3 and the annual average results are summarized in Table V.

TABLE V
AVERAGE I¹³¹ CONCENTRATIONS IN ATMOSPHERE - 1959

	$\mu\text{c I}^{131}/\text{cc}$	% MPC air-Thyroid*
Benton City, Wash.	15.4×10^{-14}	0.05
North Richland, Wash.	6.4×10^{-14}	0.02
Richland, Wash.	7.9×10^{-14}	0.03
Pasco, Wash.	13.2×10^{-14}	0.04

* Based on NCRP recommendations

Concentrations in Vegetation

Atmospheric concentrations generally result in the uptake of radionuclides by plant life. Therefore, the sampling program at Hanford includes sampling of native grasses from several locations throughout the Pacific Northwest. These locations are identified as zones as shown in Figure 15. The sampling procedure consisted of collecting 15 grams of grass at each of ten sites throughout a zone. The samples within zones were composited and analyzed by use of a gamma energy spectrometer. Radionuclides observed included Zr⁹⁵-Nb⁹⁵, Ce¹⁴¹⁺¹⁴⁴, Ru¹⁰³⁺¹⁰⁶, I¹³¹ and Ba¹⁴⁰-La¹⁴⁰. The results for all off-site zones are contained in Appendices C-4 through C-20. Figures 16, 17, 18, and 19 illustrate the variation in concentration observed during 1959. Figures 17, 18, and 19 show a similar pattern throughout the year. There is a significant difference in Figure 16 which is representative of zones located within approximately 15 miles of Hanford's chemical separations facility stacks. Measurably higher concentrations prevailed in those zones near the stacks than in the more distant zones during November and December.




FIGURE 15




FIGURE 16

Radionuclide Concentration on Native Grasses - Zone C (1959)




FIGURE 17

Radionuclide Concentration on Native Grasses - Zone O (1959)




FIGURE 18

Radionuclide Concentration on Native Grasses - Spokane &
Vicinity (1959)




FIGURE 19

Radionuclide Concentration on Native Grasses - Portland &
Vicinity (1959)

Figure 20 indicates that all sampling locations followed a similar trend throughout the year. Concentrations for Zr⁹⁵-Nb⁹⁵ only are plotted because they were consistently abundant radionuclides.

Daily measurements of I¹³¹ emissions from chemical separations effluent stacks were made and are reported in Appendix C-21. In addition, all samples of native vegetation were analyzed for I¹³¹. The criterion for release of I¹³¹ is based on uptake by vegetation and the subsequent ingestion by range animals as well as humans,⁽¹⁸⁾ rather than the breathing air MPC recommended by the NCRP. The control limit derived in the reference is $1 \times 10^{-5} \mu\text{c/g}$ of vegetation. Since this control is based upon chronic ingestion, concentrations in excess of this limit for short periods of time are relatively unimportant and do not necessarily indicate excessive exposure.

During the year there was no indication that I¹³¹ concentrations in vegetation exceeded the control limit of $1 \times 10^{-5} \mu\text{c I}^{131}/\text{g}$ in any off-site location and only rarely did the vegetation concentration exceed this level in zones within the Plant boundary. In Figure 21, the vegetation results are shown for the zone immediately surrounding one of the stacks. In this figure, the daily stack emission data have been summed, corrected for half-life, and plotted as "quantity available" according to the following equation:

$$C_0 = \sum_0^N a_t e^{-\lambda t}$$

where: C_0 = total quantity I¹³¹ available on the day of interest.

t = time in days previous to the day of interest.

a_t = the daily I¹³¹ emission on the t^{th} day previous to the day of interest.

λ = disintegration constant for I¹³¹ = 0.0845 /day.

N = earliest previous day considered, 1 November 1958 in this case.

NOTE: Because of the relatively short half-life of I¹³¹, it is not necessary to consider the amount emitted per day for more than about the 45th




FIGURE 20

Zr₉₅-Nb₉₅ Concentration on Native Grasses at Selected Sites (1959)




FIGURE 21

Comparison of I^{131} Deposition on Grasses with I^{131} Emitted in the Vicinity of a Separations Facility - 1959


previous day. However, the above equation is mathematically convenient in that the total I^{131} available from any previous day needs only to be decayed by one day and the current day's emission added to arrive at the current amount of I^{131} available.

Figure 21 indicates a tendency for grass concentrations to follow the "quantity I^{131} available" plot; however, the efficiency of "collection" of I^{131} by grass appeared to be considerably less in the summer than in the winter months.

The concentration of I^{131} in jack rabbit thyroids (Appendix C-22) followed the same pattern as the vegetation concentrations. Figure 22 compares the average jack rabbit thyroid concentrations with the average vegetation concentrations. The jack rabbits were obtained from three locations, and the data are plotted for each location. The vegetation concentration is the average concentration between dates for the area in the near vicinity of the two separations facilities' stacks. The quantity of I^{131} available was determined from the daily stack emissions; corrected for physical decay as described previously. The anomaly in thyroid and grass concentration values for October and November may be due to change in the rate of the jack rabbit's metabolism. (19)

C. Concentration of Radionuclides in Farm Products

The two farming areas located nearest downstream of the Hanford plant which use Columbia River water for irrigation are the Ringold farms and the Riverview Irrigation District. These farming areas are located about 15 and 30 miles downstream, respectively, from the nearest reactor. The farms in both areas are relatively small and so diversified that their harvests contribute only a small fraction of produce consumed in nearby communities. However, their locations with respect to the reactors should result in the maximum concentrations of radionuclides in produce for those radionuclides associated with irrigation water.



Comparison of Average I^{131} Concentrations - Jack Rabbit Thyroids and Native Grasses (1959)

FIGURE 22

The wind direction near Hanford's Separation Facilities is predominately from the northwest. Ringold is about 20 miles east and Riverview about 30 miles southeast of the chemical separations facilities, and hence are also subject to air-borne effluents. Therefore, sampling of produce was established for two additional localities which are subject to air-borne effluents but not to reactor effluent water. One location is about 20 miles south of the separations area in the vicinity of Benton City, Washington. The second location is about 40 miles southeast of the plant in the vicinity of Finley, Washington. The Yakima River is the source of irrigation water for both locations. These locations and other features of the Hanford environs are shown in Figure 1.

The foodstuffs sampled and analyzed have been categorized as to grain, vegetables, fruits, alfalfa, milk, and miscellaneous. The choice of categories and selection of samples was based on crop availability, importance to human diet, and the possibility of an unusual affinity for a particular radionuclide. For the most part, these foodstuffs were freshly harvested and were analyzed within a few days after acquisition. There was no culinary treatment given to the samples in preparation for analysis, although some depletion of radioactive contamination during washing, peeling, etc., might be expected if the produce was being prepared for human consumption.

Milk sampling consisted of samples from Ringold, Riverview and from all commercially available milk in the Richland-Pasco-Kennewick area.

Multichannel gamma energy spectrometry was used to analyze the samples for the following radionuclides: K⁴⁰, Sc⁴⁶, Cr⁵¹, Zn⁶⁵, Zr⁹⁵-Nb⁹⁵, Ru¹⁰³⁺¹⁰⁶, I¹³¹, Cs¹³⁷, Ce¹⁴¹⁺¹⁴⁴. Analysis for P³², Sr⁸⁹ and Sr⁹⁰ was accomplished through radiochemical separation and beta counting, as described in Appendix A.

The preceding radionuclides are not all of those potentially present in foodstuffs but are those which are either anticipated from Hanford's waste releases or fallout and are radiologically significant.

In tabulating the data, reference is often made to reporting limits. These reporting limits reflect some of the uncertainties of gamma energy spectra measurement of a radionuclide in the presence of other gamma emitters. Thus the reporting limit is generally higher for any particular radionuclide than the detection limit would be if the radionuclide were the only one present. The reporting limits bear no uniform mathematical relationship with detection limits but have been established through experience.

Concentrations in Milk

Small amounts of Zn⁶⁵ occurred in milk originating from areas irrigated with Columbia River water. The concentrations of Zn⁶⁵ found in milk are given in Appendix D-1 and summarized in Table VI.

TABLE VI
AVERAGE CONCENTRATIONS OF Zn⁶⁵ IN MILK

<u>Location</u>	<u>Number of Samples</u>	<u>Concentration</u> (Units of $10^{-6} \mu\text{c Zn}^{65}/\text{g}$)
Ringold	1	0.42
Riverview 1	10	< 0.05
Riverview 2	3	0.53
Riverview 3	12	1.9

The variability of occurrence of Zn⁶⁵ in milk on land irrigated with river water, as indicated by Table VI, may be influenced by the manner of irrigation of the animal's grazing food. Another possibility is the origin of the animal's drinking water which may be directly from the irrigation ditch or from a well.

Table VII shows the concentration of Zn⁶⁵ in milk over several months from a representative area and indicates little variability with season.

TABLE VII

Zn⁶⁵ IN MILK DURING THE SUMMER AND FALL OF 1959⁽¹⁷⁾

Riverview Irrigation District, Pasco, Washington
Location 3

<u>Date</u>	<u>10⁻⁶ μc/cc</u>
July 16	1.6
July 22	1.7
July 29	1.8
August 12	2.3
August 19	2.0
August 28	1.8
September 1	2.2
September 9	2.3
September 30	2.6
October 19	1.2
October 21	1.7
October 28	2.0
AVERAGE	1.93

In addition to sampling milk from local farms, milk purchased in local commercial establishments was sampled and analyzed. These results are reported in Appendix D-1 and compared with locally obtained milk in Table VIII. Small amounts of Cs¹³⁷, Sr⁸⁹, and Sr⁹⁰ were present in most milk samples.

TABLE VIII
AVERAGE CONCENTRATIONS OF SEVERAL RADIONUCLIDES IN MILK
(Units of 10^{-6} $\mu\text{c/g}$)

	No. Samples Analyzed for:	Cs ¹³⁷	No. Samples Analyzed for:	Sr ⁸⁹	Sr ⁹⁰
<u>Reporting Limits</u>		0.02		0.004	0.002
Riverview 1	10	0.044	9	0.022	0.0032
Riverview 2	3	0.025	2	0.0087	0.0036
Ringold 1	1	<0.02	1	0.0060	0.0020
Commercial A	12	0.029	7	0.0098	0.0032
Commercial B	5	0.033	1	0.0087	0.0032
Commercial C	6	0.027	4	0.0048	0.0030
Commercial D	4	0.028	3	0.0071	0.0035
Commercial E	3	0.038	2	0.021	0.0046
Commercial F	5	<0.020	2	0.010	0.0013
Commercial G	6	0.028	5	0.0079	0.0034
Commercial H	9	0.067	6	0.024	0.0087

In both local and commercial milk samples many of the radionuclides other than Cs¹³⁷ did not appear in concentrations above their respective reporting limits as shown in Appendix D-1. Such radionuclides were: Sc⁴⁶, Cr⁵¹, Zn⁶⁵, Zr⁹⁵-Nb⁹⁵, Ru¹⁰³⁺¹⁰⁶, I¹³¹, and Ce¹⁴¹⁺¹⁴⁴.

With the exception of commercial brand H, the concentrations of Cs¹³⁷, Sr⁸⁹ and Sr⁹⁰ appear quite comparable. Commercial milk H is known to include milk imported from western Washington. The higher values observed for milk H are consistent with the assumptions; the origin of these radionuclides is debris from nuclear detonations, and increased rainfall increases the amount of these radionuclides on vegetation.

The absence of detectable quantities of Zn⁶⁵ in commercial milk as indicated by the data in Appendix D-1 is probably due to the small fraction of milk contributed by the local farms to the total commercial milk supply.

Concentrations in Fresh Farm Produce

Phosphorus-32, Sc⁴⁶, Cr⁵¹ and Zn⁶⁵ are predominately reactor effluent radionuclides and as such would be expected to appear in produce from downstream farms irrigated with Columbia River water. However, the analyses performed and reported in Appendices D-3 through D-6 did not exclude the measurement of radionuclides from other sources.

Average concentrations of Zn⁶⁵ and Cr⁵¹ in produce are summarized in Tables IX and X, respectively.

TABLE IX
AVERAGE CONCENTRATIONS OF Zn⁶⁵ IN PRODUCE
AT VARIOUS LOCATIONS
(Units of 10⁻⁶ μ c/g)

	<u>Riverview</u>	<u>Ringold</u>	<u>Benton City</u>	<u>Finley</u>
Grain	1.5	2.6	< 0.1	< 0.1
Vegetables	0.18	0.50	0.12	< 0.1
Fruit	0.13	< 0.1	< 0.1	< 0.1
Alfalfa	1.3	0.68	< 0.1	< 0.1

TABLE X
AVERAGE CONCENTRATIONS OF Cr⁵¹ IN PRODUCE
AT VARIOUS LOCATIONS
(Units of 10⁻⁶ μ c/g)

	<u>Riverview</u>	<u>Ringold</u>	<u>Benton City</u>	<u>Finley</u>
Grain	0.7	< 0.5	< 0.5	< 0.5
Vegetables	0.53	0.62	< 0.5	< 0.5
Fruit	0.66	< 0.5	< 0.5	< 0.5
Alfalfa	6.6	3.4	0.64	0.58

Differences observed in results for similar kinds of produce from farms within the same district were theorized to be due to the method of irrigation. Table XI summarizes the results obtained for produce grown under the two general methods of irrigation.

TABLE XI
COMPARISON OF Zn⁶⁵ IN PRODUCE
BETWEEN DIFFERENT MODES OF IRRIGATION⁽¹⁷⁾
(Units of 10⁻⁶ μ c/g)

<u>Sample</u>	<u>Sprinkler Irrigated</u>	<u>Ditch Irrigated</u>
Alfalfa	8.9	0.88
Corn	1.4	1.6
Green Beans	1.3	0.10
Carrots	0.36	0.17
Tomatoes	0.36	0.035
Cabbage	0.18	0.28
Irrigation Water	0.072	0.070

Table XII compares the occurrences of Zn⁶⁵ and Cr⁵¹ in the Riverview district for several kinds of produces.

TABLE XII
COMPARISON OF Cr⁵¹ AND Zn⁶⁵ IN FARM PRODUCE⁽¹⁷⁾
(10⁻⁶ μ c/g)

<u>Sample</u>	<u>Cr⁵¹</u>	<u>Zn⁶⁵</u>
Red Beets*	0.13	0.16
Wheat*	0.79	4.0
Carrots*	0.29	0.17
Pasture Grass*	8.0	36.0

TABLE XII (Contd.)

<u>Sample</u>	<u>Cr⁵¹</u>	<u>Zn⁶⁵</u>
Hamburger**	< 0.22	4.5
Milk**	< 0.22	1.9
Cabbage***	0.76	0.18
Lettuce***	1.2	0.44
Boysenberries***	0.66	0.29
Pasture Grass***	6.9	3.9
Carrots***	1.0	0.36
Grapes***	0.86	0.18
Tomatoes***	0.82	0.36
Soil*** (to depth of two inches)	38.0	5.6
Irrigation Water	3.2	0.068

* Irrigated by ditch, or by flooding in the case of pasture grass and wheat.

** From cattle raised on sprinkler-irrigated pasture.

*** Sprinkler-irrigated.

Scandium-46 concentration in alfalfa from Riverview and Ringold districts averaged about $1.5 \times 10^{-7} \mu\text{c Sc}^{46}/\text{g}$ and was less than $1 \times 10^{-7} \mu\text{c/g}$ for other locations. Scandium-46 concentrations in other kinds of produce were also less than $1 \times 10^{-7} \mu\text{c/g}$.

The radionuclides; Sr⁸⁹, Sr⁹⁰, Zr⁹⁵-Nb⁹⁵, Ru¹⁰³⁺¹⁰⁶, I¹³¹, Cs¹³⁷, and Ce¹⁴¹⁺¹⁴⁴, result primarily from air-borne effluents, although they do exist in small amounts in the Columbia River water. The occurrences of these radionuclides are summarized by locality and particular kind of produce in Tables XIII through XVI.

TABLE XIII
AVERAGE CONCENTRATIONS OF SEVERAL RADIONUCLIDES
AT VARIOUS LOCATIONS FOR GRAIN

	(Units of 10^{-6} $\mu\text{c/g}$)			
	Riverview	Ringold	Benton City	Finley
Sr ⁸⁹	0.026	0.052	0.023	< 0.01
Sr ⁹⁰	0.012	< 0.006	0.020	< 0.006
Zr ⁹⁵ -Nb ⁹⁵	0.33	0.52	0.33	0.15
Ru ¹⁰³⁺¹⁰⁶	0.60	0.77	< 0.5	< 0.5
I ¹³¹	< 0.7	< 0.7	< 0.7	< 0.7
Cs ¹³⁷	< 0.05	0.083	0.057	0.059
Ce ¹⁴¹⁺¹⁴⁴	1.1	0.95	0.62	0.63

TABLE XIV
AVERAGE CONCENTRATIONS OF SEVERAL RADIONUCLIDES
AT VARIOUS LOCATIONS FOR VEGETABLES

	(Units of 10^{-6} $\mu\text{c/g}$)			
	Riverview	Ringold	Benton City	Finley
Sr ⁸⁹	0.014	< 0.01	< 0.01	0.013
Sr ⁹⁰	0.0093	< 0.006	0.0063	0.0094
Zr ⁹⁵ -Nb ⁹⁵	0.1	< 0.1	< 0.1	0.12
Ru ¹⁰³⁺¹⁰⁶	< 0.5	0.51	< 0.5	< 0.5
I ¹³¹	< 0.1	0.18	< 0.1	< 0.1
Cs ¹³⁷	0.052	< 0.05	< 0.05	< 0.05
Ce ¹⁴¹⁺¹⁴⁴	0.53	0.62	< 0.5	< 0.5

TABLE XV

AVERAGE CONCENTRATIONS OF SEVERAL RADIONUCLIDES

AT VARIOUS LOCATIONS FOR FRUIT

(Units of $10^{-6} \mu\text{c/g}$)

	Riverview	Ringold	Benton City	Finley
Sr ⁸⁹	0.61	0.63	0.011	< 0.01
Sr ⁹⁰	0.0061	0.0074	0.0074	< 0.006
Zr ⁹⁵ -Nb ⁹⁵	0.12	< 0.1	< 0.1	0.14
Ru ¹⁰³⁺¹⁰⁶	< 0.5	0.63	< 0.5	< 0.5
I ¹³¹	< 0.1	< 0.1	< 0.1	0.11
Cs ¹³⁷	< 0.05	< 0.05	< 0.05	0.57
Ce ¹⁴¹⁺¹⁴⁴	0.61	0.63	0.5	0.61

TABLE XVI

AVERAGE CONCENTRATIONS OF SEVERAL RADIONUCLIDES

AT VARIOUS LOCATIONS FOR ALFALFA

(Units of $10^{-6} \mu\text{c/g}$)

	Riverview	Ringold	Benton City	Finley
Sr ⁸⁹	0.081	0.062	0.10	0.034
Sr ⁹⁰	0.022	0.009	0.073	0.069
Zr ⁹⁵ -Nb ⁹⁵	1.5	0.17	1.7	0.4
Ru ¹⁰³⁺¹⁰⁶	0.97	0.82	1.0	1.4
I ¹³¹	0.17	0.25	0.16	0.52
Cs ¹³⁷	0.14	0.088	0.14	0.23
Ce ¹⁴¹⁺¹⁴⁴	2.6	1.3	2.4	2.7

As evidenced in Tables 13 through 16, the concentrations of radionuclides in the several kinds of produce were quite erratic. Therefore, it is difficult to distinguish between the quantity of these radionuclides originating from releases from process stacks and those originating from reactor cooling water.

Radionuclides in Miscellaneous Foodstuffs

Four samples of ground beef were purchased from local markets and analyzed for the same radionuclides discussed in the vegetable produce samples. In addition to naturally-occurring K⁴⁰, the meat was found to contain a concentration of about 3×10^{-7} μc Zr⁹⁵-Nb⁹⁵/g. In one sample only, a Cs¹³⁷ concentration of about 6×10^{-8} $\mu\text{c}/\text{g}$ was measured.

Five commercially canned baby food samples were collected and analyzed. Cesium-137 was the predominant radionuclide with a concentration of about 1×10^{-7} $\mu\text{c}/\text{g}$ in meats. The Cs¹³⁷ concentration was less than 5×10^{-8} $\mu\text{c}/\text{g}$ in the canned vegetables.

Four samples of condensed or evaporated milk were purchased and analyzed. Cesium-137 ranged from 4.1×10^{-8} to 1.7×10^{-7} $\mu\text{c}/\text{g}$. One sample indicated Ce¹⁴¹⁺¹⁴⁴ was present in the amount of 4.6×10^{-6} $\mu\text{c}/\text{g}$.

Eleven samples of dry or powdered milk were analyzed. These samples covered five different brands. The sampling period was during the last quarter of 1959. The predominant radionuclide was again Cs¹³⁷ which was present in similar amounts for all samples. The average concentration was about 4.4×10^{-7} μc Cs¹³⁷/g. In one sample, the Ce¹⁴¹⁺¹⁴⁴ concentration was 1.9×10^{-6} $\mu\text{c}/\text{g}$.

In all of the above miscellaneous foodstuffs, analysis for Sr⁹⁰ and Sr⁸⁹ was not completed.

IV. ESTIMATE OF RADIATION EXPOSURE - NONOCCUPATIONAL

Persons living in the neighborhood of the Hanford facilities receive radiation exposure from radionuclides through several paths of intake and from external radiations. In many cases it is not possible to differentiate the origins of radiation exposure among Plant operation, fallout from nuclear detonations, and natural background. However, multiple sources and paths of intake must be jointly considered in the estimate of radiation exposure.

There is a general lack of detailed information on variation of dietary habits of individuals within a population group. Because of this and other uncertainties, the radiation exposure to persons in the neighborhood of a controlled area cannot be stated precisely; although direct measurement of body burdens by whole body monitoring techniques promises improvement of the situation for some radionuclides in the future. Nevertheless, it is desirable to estimate the magnitude of exposure to man based on the data included in this report.

In summing radiation exposure from the various sources, the criteria used are those recommended by the NCRP⁽⁵⁾ for persons in the neighborhood of controlled areas.

The calculations include the recommended practice of using one-tenth of the continuous occupational limits for persons in the neighborhood of a controlled area. In addition to the assumptions inherent in the derivation of maximum permissible limits, several additional assumptions have been made as follows:

1. The persons considered resided in Pasco and obtained drinking water from the city system, used the Columbia River for swimming and boating, obtained milk from commercial sources, obtained produce from Riverview Irrigation District, consumed oysters from Willapa Bay, and obtained fish and ducks from the Columbia River downstream of the project.

2. The continuous consumption rate of foods in kg/week consisted of: milk 6, vegetables 3, fruit 3, cereal grains 1.5, oysters 0.1 and whitefish or ducks 0.1.

3. The metabolic uptake of radionuclides ingested with foodstuffs is comparable to the uptake through drinking water.

4. For foodstuffs, and where the organ of interest is the GI tract, the more restrictive limit for the radionuclide, that is soluble or insoluble form, is used to estimate the dose to the GI tract.

5. Where two radionuclides are not routinely differentiated analytically, e.g., Zr⁹⁵-Nb⁹⁵, all of the radioactive material is assumed to be the radionuclide with the more restrictive limit.

6. Radionuclides present in drinking water are soluble in body fluids.

For the purpose of this estimate, consideration was given to the consumption of varying amounts of foods as compared to the rate of water intake as used by the NCRP. It is, therefore, convenient to define the maximum permissible rate of intake (MPRI) as the maximum permissible concentration in water (MPC_w) times the rate of water intake, as defined for the standard man.

The radiation exposure of the gastrointestinal tract reaches equilibrium promptly with the intake of radionuclides under condition of chronic exposure. The recommended annual limit for GI tract dose is 1500 mrem for persons in the vicinity of controlled areas. The particular mixture of radionuclides present in the sanitary water at Pasco, as indicated by the 1959 data, results in a calculated average of ~5 per cent of the MPC_w for the GI tract or a corresponding dose of ~75 mrem per year. About 40 per cent of the GI tract dose from drinking water is from As⁷⁶, another 40 per cent from Np²³⁹, and the balance distributed among several radionuclides as illustrated previously in Figure 6.

The consumption of various foodstuffs adds to the dose to the GI tract, in mrem /yr., approximately as follows: milk 10-60, vegetables 15, fruit 15, cereal grains 25, oysters 10 and whitefish or ducks 45. Many measurements of radionuclides in milk were less than reporting limits, therefore the range given reflects alternate use of zero and reporting limit concentrations. For all other measurements the ranges are small and representative values are used. The total exposure to the GI tract from

sanitary water and locally grown foodstuffs amounted to about 250
mrem/yr.*

The dose to bone does not necessarily reach equilibrium promptly with the rate of ingestion of radionuclides. Therefore, the total fraction of the MPRI for those radionuclides which contribute to the dose to organs other than the GI tract is perhaps more illustrative than the estimated dose.

The Pasco drinking water provided < 1 per cent of the MPRI for bone, mostly from P³² and Sr⁹⁰. Milk provided another 1.5 per cent, vegetables 1.5 per cent, fruit 1 per cent, cereal grains 1 per cent, oysters < 0.5 per cent and whitefish or ducks < 5 per cent. Sr⁹⁰ is the radionuclide of major interest in all of the foregoing foodstuffs, except fish and waterfowl taken locally from the Columbia River, in which case P³² is of major concern. The percentage MPRI for bone, including all paths of intake assumed in the illustration, was ~10 per cent.

Again using the percent MPRI for radionuclides which contribute to total body exposure, a total of ~5 per cent was obtained. The distribution among the paths of intake was < 0.5 per cent from Pasco sanitary water, ~0.5 per cent from milk, ~0.5 per cent from vegetables, ~0.5 per cent from fruit, ~0.5 per cent from cereal grains, < 0.5 per cent from oysters and ~3 per cent from whitefish or ducks.

The whole body penetrating radiation exposure from external sources of Hanford origin was estimated at 6 mr/year from swimming 240 hours in the Columbia River (or from twice that number of hours of boating). Such exposure is ~1 per cent of the applicable maximum permissible limit. There may have been a small contribution to the thyroid dose as a result of I¹³¹ intake, primarily from foodstuffs. This is estimated to have been < 150 mrem per year.

* Since this estimate is based on a Pasco resident ingesting locally grown foodstuffs, the average exposure to all persons in the neighboring communities of the Hanford controlled area is less than 250 mrem/yr.

The preceding estimates fit within the framework of variation of individual dietary habits and other uncertainties. However, it is not intended that they represent boundary conditions. By comparison with relevant permissible limits for persons in the vicinity of controlled areas, the situation for Hanford during 1959 remained substantially as stated by H. M. Parker during recent Congressional Hearings.

"In summary, for man in the Hanford environs, the GI tract dose is generally the highest organ dose. Expressed as percentage of the corresponding permissible limits, GI tract dose, bone dose, and reproductive-organ dose rank about equally. The estimated representative contribution in each case is in the range of 3 per cent to 15 per cent of the limit. The corresponding maxima for exceptional cases where unusual amounts of local fish and leafy vegetables are eaten seem to be contained within the general range of say 40 per cent to 60 per cent of the limits."

V. ACKNOWLEDGEMENTS

The cooperation of many General Electric Company personnel who prepared and provided data, and reviewed this document is gratefully acknowledged. The collection of the wide variety of samples from many localities was accomplished by the Environmental Monitoring Operation under the supervision of R. C. Dozer. All routine analyses were performed by the Radiological Chemical Analysis Operation, F. E. Holt, Supervisor. Specific analyses performed by research groups and which are related to environmental measurements have been referenced in the bibliography.

The cooperation and provision of information by the United States Geological Survey Records Center, Portland, Oregon; the Pasco, Washington, City Water Department and the several state and federal agencies who operated air filter sample stations contributed substantially to the report.

BIBLIOGRAPHY

1. General Electric Co., Hanford Laboratories Staff, HAPO. Quarterly Progress Report, Research and Development Activities in the Field of Radiological Sciences, September - December 1959. Janaury 25, 1960.
2. Parker, H. M. Hearings Before the Special Subcommittee on Radiation of The Joint Committee on Atomic Energy Congress of The United States Eighty-Sixth Congress First Session on Industrial Radioactive Waste Disposal. Vol. 1, p.230. January 28-February 3, 1959.
3. Nielsen, J. M., and R. W. Perkins. The Depletion of Radioisotopes From the Columbia River by Natural Processes, HW- 52908. October, 1957. (SECRET).
4. Junkins, R. L. "Surveillance of a Municipal Water System to Assure Control of Radioisotopes - A Study of the Hanford Atomic Products Operation," 1960 Nuclear Congress, Paper No. 39. New York: 1960.
5. Maximum Permissible Amounts of Radioisotopes in the Human Body and Maximum Permissible Concentrations in Air and Water, NBS Handbook 69, June 5, 1959
6. Silker, W. B., and L. J. Kirby. Separation and Determination of Radioactive Rare Earths in Reactor Effluent Water, HW-48021. January, 1957.
7. Hungate, F. P. Biology Operation, General Electric Company, HAPO. Private Communication
8. Foster, R. F., and R. L. Junkins. Off-Project Exposure From Hanford Reactor Effluent, HW-63654. February, 1960.
9. Watson, D. G., and J. J. Davis. Concentration of Radioisotopes in Columbia River Whitefish in the Vicinity of the Hanford Atomic Products Operation, HW-48523. February, 1957. (DECLASSIFIED).
10. Davis, J. J., D. G. Watson, and C. C. Palmeter. Radiobiological Studies of the Columbia River Through December, 1955, HW-36074. November, 1956. (DECLASSIFIED).

11. Davis, J. J. et al. "Radioactive Materials in Aquatic and Terrestrial Organisms Exposed to Reactor Effluent Water," Second International Conference on the Peaceful Uses of Atomic Energy, Paper No. 393. Geneva, Switzerland: 1958.
12. Hanson, W. C., and J. J. Davis. Radioactive Contamination in Wildlife, Hanford Biology Research - Annual Report for 1958, HW-59500, pp. 138-143. January, 1959.
13. Rose, R. E. Radiological Physics Division Semiannual Report, Argonne National Laboratory, ANL-5755, pp. 53-56. January-June 1957.
14. Perkins, R. W., and J. M. Nielsen. "Zinc-65 in Foods and People," Science, 129:94-95. January 9, 1959.
15. Roesch, W. C. Radiological Physics Operation, General Electric Co., HAPO. Private Communication.
16. Bliss, C. I. "Fitting the Negative Binomial Distribution to Biological Data," (Note on the Efficient Fitting of the Negative Binomial, R. A. Fisher) Biometric, 9: 176-200. 1953.
17. Nielsen, J. M. Radiological Chemistry Operation Semiannual Report, July-December, 1959. HW-63824. January, 1960.
18. Parker, H. M. Hearings Before the Special Subcommittee on Radiation of the Joint Committee on Atomic Energy, Congress of the United States, Eighty-Sixth Congress First Session on Industrial Radioactive Waste Disposal, January 28-February 3, 1959. Vol. 1, p. 391.
19. Hanson, W. C. Biology Operation, General Electric Co., HAPO. Private Communication.

APPENDIX A

ANALYTICAL METHODS

1. Water Analyses*

All water samples are analyzed for alpha emitters, beta emitters, and selected radionuclides. Alpha emitters are extracted with diethyl ether from 9 N nitric acid. The gross alpha activity is measured with an alpha scintillation counter. Gross beta activity is determined by evaporating a sample to dryness and counting the residual salts on a gas-flow proportional beta counter.

Rare earths plus yttrium, silicon-31, iodine-131, phosphorous-32, strontium-89 and strontium-90 are measured by beta counting after chemical separation. The rare earths are isolated as a group by hydroxide, fluoride, and oxalate precipitations; silicon is precipitated as the dioxide; iodine is isolated by carbon tetrachloride extraction and precipitation as silver iodide; phosphorous by extraction of phosphomolybdic acid with butanol in diethyl ether; and strontium by successive precipitation of the nitrate and the carbonate. Yttrium-90, separated from the strontium after secular equilibrium is established, is measured to determine strontium-90. Beta decay curves are extrapolated to sampling time to determine the initial activity levels and to check separation effectiveness.

Manganese-56, zinc-69 and gallium-72 are determined by measurement of their characteristic gamma peaks with a multi-channel gamma energy spectrometer using a 3 inch x 3 inch thallium-activated sodium iodide (NaI(Th)) scintillation crystal detector. The measurements are made after the following chemical separations; manganese by precipitation as the dioxide, zinc by precipitation as the phosphate and ion exchange purification, and gallium by extraction with iso-propyl ether and precipitation as the

* L. F. Lust. Radiological Chemical Analysis Operation, General Electric Company, HAPO, Private Communication.

hydroxide. Sodium-24, neptunium-239, chromium-51 and cobalt-60 are also determined using a multi-channel gamma energy spectrometer, but are determined from a direct count of residual salts from the evaporated sample, without chemical separations. However, it may be necessary to chemically separate neptunium-239 and cobalt-60 for samples with low concentrations.

Copper-64 is determined from gamma-coincidence counting measurements of the annihilation photons produced by positron emission. Scandium-46 is measured by gamma-coincidence counting of the 0.885 Mev and 1.12 Mev photons.

Arsenic-76 is determined from the counting rate of its 2.97 Mev beta. Particles of lower energy from other beta emitters are shielded out by use of a 504 mg/cm^2 absorber.

Uranium concentrations are determined with a fluorophotometer, using standard techniques.

2. Vegetation and Produce Analyses*

Samples of native grasses (vegetation samples) are collected routinely and analyzed with a multi-channel gamma energy spectrometer for selected radionuclides. The spectrometer utilizes the 3 inch x 3 inch NaI(Th) scintillation crystal used in analyzing water samples. These analyses are conducted for 150 gram samples which have been shredded and placed in a 9-ounce glass jar. Background analysis includes the effects of the jar glass which contains minute amounts of radioactivity. There is no ashing or chemical separation performed on vegetation samples.

Farm products, including milk, are analyzed for several radionuclides including those measured in vegetation samples. Since farm produce represents a more direct means of exposure to man and because of the relatively short growing season, more sensitivity in the analytical method is desired.

* L. S. Kellogg. Radiological Chemical Analysis Operation, General Electric Company, HAPO. Private Communication.

and the phosphorous is precipitated as magnesium ammonium phosphate. After dissolving the precipitate in hydrochloric acid, ammonium citrate is again added and phosphorous is reprecipitated as magnesium ammonium phosphate.

The precipitate is dried in a $1\frac{1}{2}$ -inch stainless steel counting dish under heat lamps and counted over a period of two weeks in a gas-flow, proportional beta counter.

3. Air Sample Analyses*

Air-borne concentrations of radioactive materials and radiation dosage rates are measured by equipment located in 31 atmospheric monitoring stations, and by portable ionization chambers placed at selected field locations. Twenty-seven of the monitoring stations are within project territory, while the remainder are situated in the four perimeter towns located south and southeast of the project.

Iodine-131 scrubber samplers are operated in twelve of the monitoring stations concentrated around the separations plants. These samplers consist of a calibrated, electrically-driven vacuum pump which draws 2.0 cfm ($3.4 \text{ m}^3/\text{hr}$) of air through one liter of 0.1 normal NaOH solution. A balancing platform and siphon arrangement permits introduction of distilled water into the scrubber at a rate equal to the rate of evaporation. This water feeder helps maintain constant liquid head, air flow rate, and scrubber efficiency.

After one week of operation, the scrubber bottle is replaced and taken to the radiochemical analysis laboratory for determination of the iodine-131 content. The analytical procedure used provides for the addition of an iodine carrier and AgNO_3 to the scrubber solution, followed by filtration of the resulting silver iodide precipitate. The radiation from the I^{131} on the filter is measured by an end-window GM tube connected to a scale-of-64 scaler. Atmospheric concentrations of I^{131} are then calculated

* Soldat, J. K. Monitoring for Air-Borne Radioactive Materials at Hanford Atomic Products Operation. Presented at the Air Pollution Control Association Meeting in Los Angeles, California, June 21-26, 1959. To be published.

Increased sensitivity is achieved by using a 9-inch diameter well-type NaI(Th) scintillation crystal as the detector of a multi-channel gamma energy spectrometer. In addition, the analysis includes a determination of the radiostrontium and radiophosphorous after chemical separation. The chemical separation for radiostrontium analysis is performed in the following manner:

500 g samples of produce and 1000 g samples of milk are first dehydrated and then ashed at 500-650°C from four to six hours. The ash is dissolved in nitric acid, barium and strontium carriers are added, and the alkaline earths are precipitated as carbonates. Calcium is removed by controlled nitrate precipitation of strontium as alkaline earth metals with fuming nitric acid, washing with acetone, dissolution and reprecipitation with fuming nitric acid. The rare earths are removed from an aqueous solution of the nitrates by a Fe(OH)_3 precipitation and barium is removed as the chromate. Strontium is precipitated as the carbonate and then dried in a one-inch stainless steel counting dish to constant weight. The strontium mixture is counted for one hour in a low background (anti-coincidence) gas-flow proportional beta counter.

Strontium-90 is allowed to reach secular equilibrium with its daughter, yttrium-90, which is then extracted with buffered TTA. Yttrium-90 is counted in the same manner as the strontium mixture. The strontium-90 content is calculated from the yttrium-90 counting rate, and the strontium-89 content from the difference in counting rates of total strontium and strontium-90.

The chemical separation for radiophosphorous is performed on samples of sufficient size to yield 40-50 mg of phosphorus.

The sample is wet ashed with nitric acid. Phosphorous is precipitated from the acid solution as ammonium phosphomolybdate. This precipitate is dissolved in ammonium hydroxide, ammonium citrate is added to complex most of the remaining interfering elements,

from these counting rates by applying factors for counter calibration, chemical recovery of the I^{131} , scrubber efficiency and the volume of air sampled.

Measurements for concentrations of radioactive particulates in the atmosphere are made with 2 inch x 4 inch HV-70 filter paper. Twenty-four of these Motoaire filter samplers are operated within the project, four are operated in nearby towns, and nine others are located in remote locations scattered throughout the Pacific Northwest. The filters are changed on either a daily or a weekly schedule and then are autoradiographed using Eastman Kodak, Type-K, X-ray film. The filters are placed in direct contact with the film for one week, the filter is removed, and the film is developed. The developed film is viewed on a standard X-ray viewer and each image produced is counted as one radioactive particle. Air-borne concentrations of radioactive particles are calculated by dividing the number of images obtained per filter by the total volume (nominal 2.5 cfm) of air sampled.

External radiation dosage rates in the project environs result from radioactive materials suspended in the atmosphere, deposited on the ground, or in the river. Measurement of the low dosage rates encountered in the river is accomplished through the use of small portable ionization chambers and a novel chamber reader developed at Hanford.* Results from these chambers in use on land were seriously affected by the extreme temperatures encountered in this semi-arid region and therefore, are not reportable. The situation is currently under investigation.

4. Remarks on Units

Analytic results for concentrations are reported in $\mu\text{c/g}$ or in $\mu\text{c/cc}$ where the mass or volume refers to the sample material and not any particular chemical element. There are some results which cannot reasonably be reported in microcuries. This is due to either insufficient information regarding the particular radionuclides present in the sample and/or insufficient accuracy in the sampling method, i. e., the data reported in Appendix B-4, Alpha Emitters in Sanitary Water. Suitable isotopic analysis of the water

* Roesch, W. C., R. C. McCall and F. L. Rising. A Pulse Reading Method for Condenser Ion Chambers. Health Physics Journal 1, No. 3, pp. 340-344. December, 1958.

samples could be made thereby enabling one to convert from disintegrations per second per cc to μc per cc. However, this would add considerably to the cost of analysis. The additional cost is not justified since the 1959 results, as represented by the data in Appendix B-4, are extremely small even when compared to the MPC for the most toxic alpha emitter. Therefore these data are more useful in observing trends rather than estimating exposure.

Data which are not reported in units of microcuries for the preceding reasons are contained in the following appendices:

<u>Appendix</u>	<u>Title</u>	<u>Units</u>
B-4	Alpha Emitters in Sanitary Water - 1959	$\alpha\text{-d/s/cc}$
B-8	Beta Emitters in River Mud - 1959	$\beta\text{-d/s/g}$
B-9	Beta Emitters in Columbia River Water - 1959	$\beta\text{-d/s/cc}$
C-2	Average Beta Activity on Filters from Selected Northwest U. S. Sampling Stations - 1959	$\beta\text{-d/s/cc}$

APPENDIX B-1

CONCENTRATIONS OF SEVERAL RADIONUCLIDES IN SANITARY WATER

AT PASCO, WASHINGTON - 1959

Date	RE+Y	Na ²⁴	P ³²	Cr ⁵¹	Cu ⁶⁴	Zn ⁶⁵	As ⁷⁶	Sr ⁸⁹⁺	Sr ⁹⁰	Np ²³⁹	Zn ^{69m}
10 ⁻⁹ μ c/cc											
1-6-59	7.5	57	21	7,100	30	80	45	7.0	<0.68	4,600	NA
1-13-59	8.2	640	21	11,000	240	140	320	7.9	0.71	4,000	NA
1-20-59	15	610	15	4,900	250	13	900	4.9	<0.50	1,600	<10
1-27-59	<6.8	670	12	5,300	180	110	140	5.9	<0.50	2,400	<10
2-3-59	13	780	21	5,800	280	160	270	5.9	<0.75	2,700	16
2-10-59	9.4	400	37	5,400	120	52	180	Lost	Lost	1,800	<10
2-17-59	60	740	42	8,800	340	150	620	Lost	Lost	3,200	5.0
2-24-59	23	430	38	6,500	190	190	250	5.6	0.77	2,300	280
3-3-59	38	610	38	6,200	290	380	270	6.7	0.83	3,100	7.1
3-10-59	29	1,600	120	5,500	580	49	1,400	4.2	<0.37	2,100	<67
3-17-59	42	570	27	4,600	450	66	230	4.5	0.53	1,900	<36
3-24-59	60	1,200	<6.4	3,700	270	240	57	4.8	<0.48	2,100	25
3-31-59	16	790	12	5,500	170	130	95	5.0	<0.47	1,600	7.5
4-7-59	44	2,000	29	5,400	670	170	350	4.9	<0.38	2,700	45
4-14-59	43	1,500	12	4,500	590	210	190	4.5	Lost	4,100	28
4-21-59	31	1,600	18	4,200	650	150	320	3.6	<0.43	2,100	61
4-28-59	18	250	7.1	2,300	250	100	150	6.5	0.66	990	<56
5-5-59	18	370	<6.7	2,200	420	54	170	7.0	<0.34	1,500	<22
5-12-59	9.0	640	<3.7	1,300	330	73	850	3.1	<0.39	730	42
5-19-59	16	310	5.9	2,300	310	95	70	3.6	<0.91	1,100	22
5-26-59	25	1,000	7.8	2,100	520	73	260	3.5	<0.76	1,200	56
6-2-59	41	1,100	13	1,300	690	61	99	3.1	<0.41	790	<42
6-9-59	23	600	7.2	1,400	450	58	110	2.3	<0.38	600	28
6-16-59	74	330	<3.6	1,100	770	28	280	1.6	<0.36	530	<35
6-23-59	22	350	<6.6	1,000	380	25	58	2.3	<0.40	500	<99
6-30-59	14	130	<3.6	730	170	26	63	2.1	0.62	630	<16
7-7-59	8.7	250	<3.6	1,100	240	23	69	2.6	<0.67	880	<41
7-14-59	18	230	<6.7	1,300	240	21	52	1.7	0.60	520	<16
7-21-59	24	130	<6.6	1,100	340	24	70	4.2	0.57	590	<22

NA = No Analysis

APPENDIX B-1 (contd.)

Date	RE+Y	Na ²⁴	P ³²	Cr ⁵¹	Cu ⁶⁴	Zn ⁶⁵	As ⁷⁶	Sr ⁹⁰	Sr ⁹⁰⁺	Np ²³⁹	Zn ^{69m}
7-28-59	30	210	<6.2	1,300	310	240	470	2.9	<0.62	490	<14
8-4-59	34	660	14	1,800	510	32	300	3.4	0.60	650	<18
8-11-59	53	580	21	2,100	690	38	260	3.7	0.80	740	<15
8-18-59	52	1,000	28	3,900	750	48	480	4.9	<0.73	1,900	34
8-25-59	45	1,100	23	3,600	710	37	450	5.1	1.3	1,600	<33
9-1-59	26	550	23	4,000	320	46	320	5.3	1.1	1,400	<20
9-9-59	30	1,100	18	4,800	640	32	380	4.3	<0.94	2,100	<32
9-15-59	19	720	25	3,000	490	32	350	4.2	0.66	1,300	28
9-22-59	24	320	23	2,500	120	58	140	4.6	0.61	850	32
9-29-59	12	360	28	2,800	120	37	280	6.0	0.88	1,300	<15
10-6-59	44	540	65	3,000	350	60	670	4.1	0.75	1,300	<19
10-13-59	28	250	95	3,800	120	49	560	5.0	0.81	1,400	<18
10-20-59	57	550	100	3,900	420	79	940	6.0	<0.74	1,800	<21
10-27-59	56	480	120	1,900	330	100	900	5.3	0.72	1,100	<16
11-3-59	50	240	96	3,200	150	61	470	3.0	0.61	1,100	<34
11-10-59	7.4	84	8.7	1,000	76	52	250	2.6	<0.63	120	<14
11-17-59	15	30	86	3,300	31	36	49	3.8	0.70	210	<13
11-24-59	59	690	140	7,000	450	99	430	6.3	0.73	2,700	<20
12-1-59	16	81	31	3,900	59	49	130	5.2	0.53	1,500	<16
12-8-59	29	230	33	4,800	220	90	190	3.8	0.74	1,500	<45
12-15-59	130	26	130	4,200	150	60	310	5.9	0.82	1,400	<15
12-22-59	42	54	49	4,600	87	150	970	6.9	0.60	1,400	<34
12-29-59	89	170	140	4,900	150	53	480	9.7	0.51	2,300	<47

APPENDIX B-2

CONCENTRATIONS OF SEVERAL RADIONUCLIDES IN SANITARY WATER

AT KENNEWICK, WASHINGTON - 1959

Date	RE+Y	Na ²⁴	P ³²	Cr ⁵¹	Cu ⁶⁴	Zn ⁶⁵	As ⁷⁶	Sr ⁸⁹⁺	Sr ⁹⁰	Np ²³⁹	Zn ^{69 m}
1-13-59	16	210	53	4,700	200	36	480	2.1	< 0.55	840	NA
1-20-59	22	330	40	2,600	180	13	290	1.3	< 0.58	580	< 10
1-27-59	21	180	29	3,100	190	64	270	1.1	< 0.51	520	22
2-3-59	25	86	30	2,900	230	22	330	< 1.3	Lost	540	37
2-17-59	93	610	110	5,100	650	110	850	2.8	0.45	1,700	75
2-24-59	39	310	46	3,100	310	53	200	1.7	< 0.56	610	47
3-3-59	35	140	33	1,800	130	18	270	1.5	< 0.40	300	< 32
3-10-59	38	120	32	2,800	240	34	180	1.3	< 0.38	560	< 45
3-17-59	40	120	28	2,900	250	25	130	1.0	< 0.44	510	< 23
3-24-59	50	150	30	2,900	240	39	170	< 0.9	< 0.50	710	< 50
3-31-59	28	110	22	3,000	160	38	83	1.0	< 0.38	370	< 61
4-7-59	36	340	41	2,900	330	41	200	1.3	< 0.53	550	29
4-14-59	20	160	35	2,800	300	52	300	1.2	< 0.41	580	72
4-21-59	48	150	38	3,100	400	50	230	1.0	< 0.33	490	< 65
4-28-59	19	98	15	< 1,600	130	22	70	1.1	< 0.38	< 350	< 33
5-5-59	23	61	13	< 1,600	180	20	48	1.3	< 0.30	< 310	< 36
5-12-59	23	95	15	920	250	29	120	1.8	< 0.38	310	< 62
5-19-59	13	75	9.4	1,100	150	30	97	< 0.88	< 0.68	280	< 15
5-26-59	28	260	14	1,500	330	31	130	1.2	< 0.86	250	< 43
6-2-59	46	420	15	1,100	330	50	130	1.3	< 0.39	300	< 100
6-9-59	14	55	5.4	940	170	32	54	1.0	< 0.62	260	< 34
6-16-59	25	80	8.1	730	230	18	25	0.70	< 0.41	250	< 13
6-23-59	32	240	7.3	900	310	14	73	< 1.1	< 0.39	260	< 38
6-30-59	19	130	6.7	790	240	7.5	86	1.0	< 0.41	470	< 16
7-7-59	12	78	< 3.6	760	150	10	36	< 0.51	< 0.68	190	< 12
7-14-59	36	160	< 12	1,400	310	9.0	170	1.2	0.37	300	< 15
7-21-59	33	99	< 6.1	1,300	260	< 20	120	1.8	< 0.41	340	< 28
7-28-59	16	71	< 6.6	1,200	170	< 18	440	< 0.74	< 0.59	140	< 15

-67-

HW-64371

NA = No Analysis

APPENDIX B-2 (contd.)

Date	RE+Y	Na ²⁴	P ³²	Cr ⁵¹	Cu ⁶⁴	Zn ⁶⁵	As ⁷⁶	Sr ⁹⁰⁺	Sr ⁹⁰	Np ²³⁹	Zn ^{69m}
8-4-59	24	310	13	1,800	490	<20	210	1.3	<0.45	270	<13
8-11-59	22	230	10	1,700	340	<22	130	1.5	<0.38	230	<16
8-18-59	9.8	260	10	2,600	310	24	200	1.1	<0.69	800	<41
8-25-59	26	370	15	2,500	370	19	190	1.9	<0.70	420	<13
9-1-59	6.9	63	6.4	2,100	110	<20	54	<0.89	<0.75	120	<15
9-9-59	19	35	15	2,800	380	25	260	1.6	<0.63	490	<56
9-15-59	5.2	63	6.7	1,900	110	<18	20	<1.1	<0.59	98	<15
9-22-59	11	90	<6.6	1,500	82	<20	60	<1.1	0.38	110	<17
9-29-59	5.2	190	7.4	1,800	110	25	85	<0.90	<0.32	170	<18
10-6-59	9.6	160	10	2,000	170	<19	170	<1.1	<0.41	220	<16
10-13-59	7.5	120	<5.4	2,500	140	<19	120	<1.0	<0.38	200	<17
10-20-59	7.8	92	<6.5	2,200	140	<18	110	<1.1	<0.74	180	<12
10-27-59	7.1	55	<6.6	530	130	16	87	<0.96	<0.34	20	<12
11-3-59	10	82	8.7	1,900	49	<19	18	<0.62	<0.41	30	<14
11-10-59	9.3	59	<9.7	1,700	110	<20	82	<1.0	<1.4	100	<13
11-17-59	11	170	11	2,600	140	27	85	<1.2	<0.41	290	<20
11-24-59	9.0	130	13	3,300	140	<20	110	<1.2	<0.42	320	<32
12-1-59	7.0	6.6	<6.5	2,100	75	<20	<85	<0.99	<0.38	100	<24
12-8-59	14	14	11	2,000	150	<20	<80	<1.0	<0.73	190	<18
12-15-59	18	18	12	2,000	98	20	<88	<1.1	<0.39	240	<23
12-22-59	12	14	11	2,300	120	13	160	1.0	<0.57	230	<52

APPENDIX B-3

CONCENTRATIONS OF SEVERAL RADIONUCLIDES IN COLUMBIA RIVER WATER

AT PASCO, WASHINGTON - 1959

Date	RE+Y	Na ²⁴	P ³²	Cr ⁵¹	Cu ⁶⁴	Zn ⁶⁵	As ⁷⁶	Sr ⁸⁹⁺	Sr ⁹⁰	Np ²³⁹	Zn ⁶⁵ min
1-6-59	120	1,000	430	7,800	1,400	410	5,000	7.6	0.86	5,900	NA
1-13-59	200	1,800	500	11,000	2,700	510	5,400	10	0.63	6,500	NA
1-20-59	150	1,300	140	6,000	1,800	260	2,500	15	1.5	3,600	95
1-27-59	130	530	360	2,600	560	140	1,300	4.8	0.50	1,800	44
2-3-59	390	1,800	310	6,000	2,600	310	3,600	5.5	0.77	4,100	170
2-10-59	17	670	320	4,600	350	170	410	Lost	Lost	2,900	23
2-17-59	190	1,300	410	11,000	470	180	W ^b	2,600	7.9	0.56	4,000
2-24-59	320	1,100	320	6,300	1,500	450	2,200	9.6	0.94	3,900	41
3-3-59	310	910	250	6,400	1,100	420	1,800	6.9	0.88	3,400	220
3-10-59	470	1,900	320	5,8	5,300	2,600	410	2,300	5.5	3,500	400
3-17-59	390	1,200	230	6,000	1,500	350	1,100	5.3	0.95	2,800	160
3-24-59	360	1,400	250	4,400	1,400	380	1,300	4.7	< 0.49	2,500	180
3-31-59	240	870	200	5,400	890	230	890	5.5	0.41	2,300	79
4-7-59	460	2,600	300	6,900	2,500	390	2,000	5.2	0.46	3,200	290
4-14-59	450	2,600	230	5,500	2,700	400	412	2,200	5.8	? 3,000	320
4-21-59	370	1,800	220	5,100	2,600	330	2,300	4.5	0.36	2,600	330
4-28-59	140	720	110	2,500	930	220	350	5.3	0.45	1,100	120
5-5-59	300	1,200	110	3,000	3,000	190	670	6.6	0.36	1,500	210
5-12-59	130	730	62	1,500	1,000	180	640	4.3	< 0.50	880	480
5-19-59	170	1,200	87	2,400	2,200	180	1,200	2.7	0.96	1,300	130
5-26-59	210	1,200	64	2,600	2,200	150	950	2.7	< 0.95	1,300	170
6-2-59	150	800	39	1,600	1,400	150	590	3.1	0.45	750	110
6-9-59	160	800	32	1,300	1,300	130	470	2.4	0.68	740	120
6-16-59	170	690	18	1,200	1,500	100	400	1.7	0.19	580	87
6-23-59	140	370	25	970	1,400	86	410	5.4	1.1	580	24
6-30-59	130	540	21	800	1,100	67	330	2.2	< 0.34	690	54
7-7-59	170	700	22	1,500	1,400	91	490	2.3	< 0.67	760	72
7-14-59	210	340	20	1,700	1,200	55	360	1.9	< 0.37	680	43
7-21-59	100	260	21	1,000	980	59	250	3.0	< 0.40	600	22
7-28-59	130	420	24	2,800	1,100	69	300	2.6	< 0.57	620	49

-69-

HW-64371

NA = No Analysis

APPENDIX B-3 (contd.)

Date	RE+Y	Na ²⁴	P ³²	Cr ⁵¹	Cu ⁶⁴	Zn ⁶⁵	As ⁷⁶	Sr ⁸⁹⁺	Sr ⁹⁰	Np ²³⁹	Zn ^{69m}
8-4-59	200	930	43	2,000	1,700	90	930	4.4	0.57	810	59
8-11-59	190	1,400	48	2,600	2,100	120	820	4.4	0.80	1,100	66
8-18-59	140	1,300	60	4,000	2,200	130	1,200	4.6	1.0	1,800	81
8-25-59	190	1,500	78	3,500	1,800	160	1,100	3.2	0.90	1,700	61
9-1-59	69	1,200	57	3,800	1,700	120	1,300	5.5	1.0	1,800	22
9-8-59	140	1,400	83	4,700	2,000	120	1,400	4.6	< 0.65	2,300	16
9-15-59	72	1,300	85	3,400	1,800	130	211,100	4.7	0.68	1,700	25
9-22-59	170	1,400	94	2,700	2,000	150	1,400	4.4	0.58	1,600	40
9-29-59	58	1,100	140	4,000	1,700	120	1,300	5.9	0.79	2,100	18
10-6-59	130	1,300	100	3,400	1,800	160	1,300	3.9	0.81	1,800	34
10-13-59	110	710	140	3,500	910	110	2181,200	5.8	0.70	1,900	12
10-20-59	210	1,500	180	4,300	2,400	160	1,800	6.2	< 1.0	2,500	27
10-27-59	170	1,100	170	4,300	1,700	180	1,300	4.4	0.64	1,900	26
11-3-59	140	740	110	3,100	970	140	960	4.5	0.71	1,900	19
11-10-59	190	1,500	150	314,600	2,300	190	342,400	4.2	0.65	2,600	40
11-17-59	380	1,800	180	6,100	2,800	250	32,000	6.7	0.71	3,800	80
11-24-59	410	2,100	280	8,700	360	250	2,700	6.7	0.87	4,300	68
12-1-59	140	290	99	2,800	820	150	510	4.1	0.76	1,300	28
12-8-59	180	1,300	200	5,800	2,400	270	1,200	4.4	1.1	3,100	36
12-15-59	380	1,100	230	5,600	1,300	250	311,300	7.4	0.72	3,300	33
12-22-59	220	570	180	5,100	940	190	2,500	11	0.83	2,500	81
12-29-59	290	940	230	5,700	1,300	220	2,300	9.6	0.61	3,000	100

3811
37642
11
3317.6
3237.7

962 693

Flu

APPENDIX B-4

ALPHA EMISSIONS IN SANITARY WATER - 1959
Disintegrations per Second per Cubic Centimeter

<u>Location*</u>	<u>January</u>	<u>February</u>	<u>March</u>	<u>April</u>	<u>May</u>	<u>June</u>	<u>July</u>	<u>August</u>	<u>September</u>	<u>October</u>	<u>November</u>	<u>December</u>
Benton City Water Co.	0.00033	0.00017	0.00038	0.00035	0.00043	0.00058	<0.00033	<0.0001	0.00033	0.00019	NA	NA
Sacajawea Park	0.00036	0.00026	0.00032	0.00032	0.00028	0.00027	0.00045	<0.0001	0.00037	0.00018	0.00031	NA

* 19 other locations, off project, were analyzed routinely for alpha emitters in sanitary water. All results for these locations were < 5×10^{-4} d/s per cc.

** No Analysis

APPENDIX B-5

URANIUM CONCENTRATIONS IN SANITARY WATER - 1959

<u>Location*</u>	<u>January</u>	<u>February</u>	<u>March</u>	<u>April</u>	<u>May</u>	<u>June</u>	<u>July</u>	<u>August</u>	<u>September</u>	<u>October</u>	<u>November</u>	<u>December</u>
Benton City Water Co.	2.4	2.9	16	11	8.2	19	24	7.1	11	10	-	NA
100-F Area (On Project)	---	---	---	---	2.0	2.2	-	-	-	-	-	6.6
Pasco-Composite Sample	2.3	-	-	-	-	-	-	-	-	-	-	NA
Sacajawea Park	1.2	10	17	15	16	7.7	16	17	12	15	11	NA
McNary Dam	NA***	-	-	-	-	3.3	2.7	-	-	-	-	NA
Patterson Store	NA	7.1	6.3	4.5	2.7	9.9	-	4.5	-	-	6.9	NA

* 10 other locations, off project, were analyzed routinely for uranium concentration in sanitary water. All results for these locations were less than detection limit.

** Uranium concentrations below the reporting limit are indicated by a dash (-). The reporting limit prior to September 1959 was $2 \times 10^{-9} \mu\text{c}/\text{cc}$; from September 1959 to date the reporting limit is $6 \times 10^{-9} \mu\text{c}/\text{cc}$.

*** No Analysis.

APPENDIX B-6

CONCENTRATION OF P³² IN SELECTED ORGANISMS FROM THE COLUMBIA RIVER - 1959

Units - 10⁻³ μ c/g (wet)

Date	Organism	Location	Number of Samples	Average	Maximum
1/4	River Duck	Near Pasco	1	0.056	-
1/9	Minnows, Whole Fish	River near Reactors	10	2.3	3.0
1/20	River Duck	River near Reactors	1	0.19	-
1/29	River Ducks	River Ringold Area	4	0.039	0.056
1/29	Diving Ducks	River Ringold Area	4	1.6	2.9
2/2	Minnows, Whole Fish	River Priest Rapids Area	10	0.00062	0.026
2/2	Whitefish Flesh	River Priest Rapids Area	3	0.22	0.47
2/3	Whitefish Flesh	River near Reactors	3	0.11	0.12
2/5	Minnows, Whole Fish	River Ringold Area	10	1.8	1.9
3/12	Minnows, Whole Fish	River Ringold Area	11	1.1	1.5
4/13	Minnows, Whole Fish	River Ringold Area	24	0.87	1.5
5/13	Minnows, Whole Fish	River Ringold Area	16	0.91	1.9
6/10	Minnows, Whole Fish	River Ringold Area	10	1.1	1.4
7/6	Minnows, Whole Fish	River Ringold Area	10	2.4	3.4
8/3	Minnows, Whole Fish	River Ringold Area	9	10	12
8/31	Minnows, Whole Fish	River Ringold Area	10	10	14
10/12	Minnows, Whole Fish	River Ringold Area	10	17	21
10/15	River Ducks	River Ringold Area	9	0.11	0.49
10/29	Whitefish Flesh	River Ringold Area	1	0.22	-
10/30	Minnows, Whole Fish	River Priest Rapids Area	2	0.040	0.049
10/30	Whitefish Flesh	River Priest Rapids Area	13	0.67	3.5
11/13	Whitefish Flesh	River near Reactors	12	1.4	3.9
12/8	Whitefish Flesh	River near Reactors	14	0.58	1.3
12/9	Whitefish Flesh	River Ringold Area	4	0.40	0.57
12/9	Minnows, Whole Fish	River near Reactors	10	3.3	3.8
12/10	Whitefish Flesh	River Priest Rapids Area	5	0.079	0.25

APPENDIX B-7

CONCENTRATIONS OF SEVERAL RADIONUCLIDES IN
COLUMBIA RIVER WATER AT VANCOUVER, WASHINGTON - 1959

(10^{-9} $\mu\text{c}/\text{cc}$)

<u>Date</u>	<u>P³²</u>	<u>Cr⁵¹</u>	<u>Zn⁶⁵</u>	<u>Np²³⁹</u>
6-23-59	14	710	41	300*
7-28-59	5.3	940	< 20	94
8-25-59	9.8	2,300	39	87
9-29-59	19	1,800	< 19	120
10-19-59	37	Lost	--	--
11-3-59	39	2,300	< 19	160
12-1-59	80	4,000	55	400
12-17-59	43	2,800	67	93

* Questionable analytical result.

APPENDIX B-8

BETA EMITTERS IN RIVER MUD - 1959

Disintegrations per Second per Gram

<u>Location</u>	<u>January</u>	<u>February</u>	<u>March</u>	<u>April</u>	<u>May</u>	<u>June</u>	<u>July</u>	<u>August</u>	<u>September</u>	<u>October</u>	<u>November</u>	<u>December</u>
<u>Columbia River</u>												
Wills Ranch (Upstream from Project)	0.54	0.79	0.74	0.70	0.62	0.61	0.81	0.98	0.74	0.45	0.52	NA*
Hanford (old townsite)	3.2	2.5	1.3	3.5	1.2	1.2	2.0	2.2	3.3	4.4	4.5	NA
Below 300 Area (Downstream from Project)	3.2	1.9	1.7	3.6	2.5	1.4	2.1	2.1	3.2	2.3	3.0	NA
Byers Landing	1.3	2.0	2.2	1.4	0.72	0.75	0.82	1.9	1.2	1.9	2.4	NA
Richland	1.6	2.8	1.2	1.3	1.9	0.65	1.2	1.6	1.8	1.5	2.8	NA
Pasco-Kennewick Bridge-South End	5.0	1.6	1.6	1.3	0.68	0.86	0.88	2.1	1.3	1.4	0.84	NA
Sacajawea Park	1.8	2.1	2.2	1.3	1.6	0.94	0.70	1.1	1.1	1.0	1.6	NA
McNary - Below Dam	0.62	0.91	0.94	0.74	0.64	0.67	1.1	0.72	0.82	1.4	1.1	NA
Paterson	0.94	0.95	1.0	0.95	0.61	1.0	0.86	1.2	0.81	1.2	0.96	NA
<u>Other Muds</u>												
Snake River (Near Mouth)	0.49	0.86	0.85	0.69	0.52	0.58	0.58	0.86	0.57	0.53	0.84	NA

*NA = No Analysis

APPENDIX B-9BETA EMITTERS IN COLUMBIA RIVER WATER - 1959

Location*	Disintegrations per Second per Cubic Centimeter											
	January	February	March	April	May	June	July	August	September	October	November	December
Wills Ranch (Upstream from Project)	<0.0020	<0.0020	0.0034	0.00092	<0.00050	0.00074	0.00067	0.00059	<0.00050	0.00056	<0.00050	NA**
Harford (Old Townsite)	3.5	2.8	2.8	2.7	1.3	0.55	0.55	1.2	1.4	2.0	1.9	2.3
Richland Dock	0.67	0.43	1.2	1.0	0.54	0.21	0.26	0.42	0.51	0.76	0.79	1.0
Pasco Filter Plant Inlet	1.2	0.70	0.54	0.55	0.45	0.15	0.12	0.21	0.19	0.50	0.66	NA
Pasco-Kennett Bridge-North End	1.2	0.72	0.82	0.59	0.31	0.16	0.15	0.32	0.34	0.49	0.46	NA
Sacajawea Park	0.82	0.46	0.63	0.43	0.24	0.12	0.10	0.18	0.24	0.30	0.30	0.48
McNary - Below Dam	0.14	0.028	0.11	0.064	0.066	0.037	0.033	0.045	0.030	0.062	0.046	0.056
Paterson Ferry-Oregon	0.17	0.096	0.093	0.056	0.087	0.038	0.042	0.030	0.043	0.067	0.047	0.060
Hood River, Oregon	0.072	0.016	0.035	0.025	0.037	0.036	0.018	0.018	0.016	0.021	0.018	0.037
Vancouver, Washington	0.050	0.025	0.027	0.017	0.023	0.016	NA	0.0068	0.0089	0.013	0.016	0.026

* 15 other locations, off project, were analyzed routinely for beta emitters in Columbia River water. Results for these locations were < 1×10^{-4} d/s per cc.

** No Analysis.

APPENDIX B-10URANIUM CONCENTRATIONS IN RIVER WATER - 1959

Location**	10 ⁻⁹ μ c/cc											
	January	February	March	April	May	June	July	August	September	October	November	December
Columbia River												
Wills Ranch (Upstream from Project)	-*	3.1	-	3.5	2.2	2.6	3.8	2.5	-	-	-	NA
Harford (Old Townsite)	-	2.2	3.7	-	2.3	2.3	2.2	-	-	-	-	NA
Richland Dock	-	-	3.0	-	-	-	-	-	-	-	-	NA
Pasco-Kennett Bridge-North End	3.4	-	-	-	4.3	-	-	-	-	6.4	-	NA
McNary - Below Dam	NA***	3.1	3.6	-	3.2	-	2.11	-	-	-	12.1	NA
Paterson Ferry-Oregon	2.5	2.7	-	-	-	2.0	-	-	-	-	-	NA
Hood River, Oregon	2.5	-	-	-	2.6	2.4	3.1	-	-	-	-	NA
Vancouver, Washington	3.2	5.9	-	3.6	3.3	2.6	NA	2.4	2.4	-	-	NA
Other Waters												
Snake River-Near Mouth	4.1	-	-	2.0	2.5	4.0	-	2.4	-	-	9.1	NA

* Uranium concentrations below the reporting limit are indicated by a dash (-). The reporting limit prior to September 1959 was 7×10^{-5} d/s per cc; from September 1959 to date the reporting limit is 2×10^{-4} d/s per cc.

** 8 other locations, off project were analyzed routinely for uranium concentrations in river water. Results for these locations were below detection limit.

*** No Analysis.

APPENDIX B-11

 ESTIMATED RATE OF TRANSPORT FOR SEVERAL RADIONUCLIDES IN COLUMBIA RIVER WATER
 PASSING PASCO, WASHINGTON - 1959

Date	RE+Y	Na ²⁴	P ³²	Cr ⁵¹	Cu ⁶⁴	Zn ⁶⁵	Sr ⁸⁹⁺ Sr ⁹⁰	Sr ⁹⁰	Np ²³⁹	Zn ^{69m}
1-6-59	21	180	76	1,400	250	72	900	1.3	0.15	NA
1-13-59	40	360	100	2,200	550	100	1,100	2.0	0.13	NA
1-20-59	36	310	34	1,400	430	62	600	3.6	0.36	23
1-27-59	30	120	84	600	130	33	300	1.1	0.12	420
2-3-59	99	460	78	1,500	660	78	910	1.4	0.19	100
2-10-59	3.3	120	6.3	900	69	33	80	Lost	Lost	44
2-17-59	40	270	86	2,300	98	38	540	1.7	0.12	570
2-24-59	76	260	76	1,500	360	110	520	2.3	0.22	840
3-3-59	75	220	60	1,500	270	100	430	1.7	0.21	820
3-10-59	120	490	82	1,400	670	110	590	1.4	0.12	900
3-17-59	99	300	58	1,500	380	89	280	1.3	0.24	710
3-24-59	91	360	64	1,100	360	97	330	1.2	<0.12	41
3-31-59	57	210	48	1,300	210	55	210	1.3	0.098	640
4-7-59	120	690	79	1,800	660	100	530	1.4	0.12	550
4-14-59	130	740	66	1,600	770	110	630	1.7	?	850
4-21-59	120	580	71	1,700	840	110	740	1.5	0.12	77
4-28-59	56	290	44	1,000	370	89	140	2.1	0.18	440
5-5-59	160	640	59	1,600	1,100	100	360	3.5	0.19	800
5-12-59	76	430	36	870	580	100	370	2.5	<0.29	510
5-19-59	100	710	51	1,400	1,300	110	710	1.6	0.57	280
5-26-59	150	860	46	1,900	1,600	110	680	1.9	<0.68	77
6-2-59	120	660	32	1,300	1,200	120	490	2.6	0.37	120
6-9-59	140	710	28	1,100	1,100	110	410	2.1	0.60	91
6-16-59	160	640	17	1,100	1,400	92	370	1.6	0.18	650
6-23-59	150	380	26	1,000	1,500	89	430	5.6	1.1	540
6-30-59	140	560	22	830	1,100	70	340	2.3	<0.35	80
7-7-59	150	600	19	1,300	1,200	79	420	2.0	<0.58	56
7-14-59	150	400	14	1,200	860	40	260	1.4	<0.27	62
7-21-59	68	180	14	680	660	40	170	2.0	<0.27	31

NA = No Analysis

APPENDIX B-11 (contd.)

Date	RE+Y	Na ²⁴	P ³²	Cr ⁵¹	Cu ⁶⁴	Zn ⁶⁵	As ⁷⁶	Sr ⁸⁹⁺	Sr ⁹⁰	Np ²³⁹	Zn ^{69m}
7-28-59	79	260	15	1,700	670	40	490	1.6	< 0.35	380	30
8-4-59	90	420	19	1,900	760	40	420	2.0	0.26	360	26
8-11-59	69	510	18	950	770	44	300	1.6	0.29	400	24
8-18-59	41	380	18	1,200	650	38	350	1.4	0.29	530	24
8-25-59	48	380	20	890	460	41	280	0.81	0.23	430	15
9-1-59	17	300	14	940	420	30	320	1.4	0.25	440	5.4
9-8-59	34	340	20	1,100	480	29	340	1.1	< 0.16	560	3.9
9-15-59	20	360	24	950	500	36	310	1.3	0.19	470	7.0
9-22-59	62	510	34	990	730	55	510	1.6	0.21	590	15
9-29-59	20	380	49	1,400	590	42	450	2.0	0.27	730	6.2
10-6-59	36	360	28	940	500	44	360	1.1	0.22	500	9.4
10-13-59	29	190	37	930	240	29	320	1.5	0.19	510	3.2
10-20-59	64	450	55	1,300	730	48	550	1.9	< 0.30	760	8.2
10-27-59	56	360	56	1,400	790	53	430	1.5	0.21	630	8.6
11-3-59	43	230	34	950	300	43	300	1.4	0.22	590	5.9
11-10-59	56	440	44	1,300	670	56	410	1.2	0.19	760	12
11-17-59	85	400	40	1,400	630	56	450	1.5	0.16	850	18
11-24-59	110	590	78	2,400	100	70	750	1.9	0.24	1,200	19
12-1-59	48	99	34	960	280	51	170	1.4	0.26	450	9.6
12-8-59	42	300	46	1,300	550	62	280	1.0	0.25	720	8.3
12-15-59	85	250	51	1,200	290	56	290	1.7	0.16	740	7.4
12-22-59	52	140	43	1,200	220	45	590	2.6	0.20	590	19
12-29-59	59	190	47	1,200	260	44	470	1.9	0.12	610	20

APPENDIX B-12

ESTIMATED RATE OF TRANSPORT FOR SEVERAL RADIONUCLIDES
IN COLUMBIA RIVER WATER PASSING VANCOUVER, WASHINGTON

1959

curies/day

<u>Date</u>	<u>P³²</u>	<u>Cr⁵¹</u>	<u>Zn⁶⁵</u>	<u>Np²³⁹</u>
6-23-59	19	960	55	400*
7-28-59	3.8	670	< 14	67
8-25-59	3.2	750	13	28
9-29-59	8.8	830	< 9	55
10-19-59	16	Lost	--	--
11-3-59	17	1,000	< 8	70
12-1-59	39	2,000	27	200
12-17-59	16	1,000	25	34

* Questionable analytical result

APPENDIX B-13

CONCENTRATIONS OF SEVERAL RADIONUCLIDES IN LOCALLY PURCHASED OYSTERS* - 1959

$10^{-6} \mu\text{c/g}$									
<u>Code</u>	<u>Type</u>	<u>Location</u>	<u>Date</u>	<u>Sc⁴⁶</u>	<u>K⁴⁰</u>	<u>Zn⁶⁵</u>	<u>Zr⁹⁵-Nb⁹⁵</u>	<u>Cs¹³⁷</u>	<u>Ru¹⁰³⁺¹⁰⁶</u>
<u>Reporting Limits</u>				0. 1	0. 3	0. 1	0. 1	0. 05	0. 5
<u>Oysters</u>									
401	Freshly Shelled	Wash. Coast	11/25	-**	2. 8	67	-	-	-
401	Shells of above		11/25	-	0. 41	1. 4	1. 4	-	-
402	Brand A	Wash. Coast	11/25	-	2. 1	47	-	-	-
409	Brand A	Wash. Coast	12/7	-	1. 8	39	-	-	-
410	Brand B	Wash. Coast (?)	12/7	0. 16	2. 2	53	-	-	2. 7
									1. 6

* Strontium-89 and strontium-90 were less than 0. 004 and 0. 002, respectively in oysters.

** Results less than reporting limit are indicated by a (-).

HW-64371

APPENDIX B-14

Zn⁶⁵ IN LOCALLY PURCHASED OYSTERS - 1959*

<u>Sample No.</u>	<u>Packing</u>	<u>Origin</u>	<u>Zn⁶⁵ 10⁻⁶ μc/g</u>
1	Fresh	West Coast	63
2**	Fresh	Willapa Bay, Wash.	56
3	Fresh	South Bend, Wash.	38
4	Canned	West Coast	15
5	Canned Oyster Stew	Seattle, Wash.	4.1
6	Canned	Gulf of Mexico	< 0.14
7	Canned	New Orleans, La.	0.30
8	Canned	Biloxi, Mississippi	< 0.14
9	Fresh	Port Norris, N. J.	0.18
10	Canned	Japan	0.18
11	Canned	Japan	< 0.22
12	Canned	Japan	< 0.14
13	Canned	Japan	< 0.22

* Data supplied by Radiological Chemistry Operation

** Sampled September 5, 1959, from Willapa Bay near Bay Center, Washington.

APPENDIX B-15

Zn⁶⁵ IN COMMON MARINE FOOD ORGANISMS*

(Specimens collected April 1959)

<u>Location</u>	<u>Sample Type</u>	$10^{-6} \mu\text{c/g (Wet)}$
Coos Bay, Oregon	Burrowing Clams (soft parts)	0.45
	White Perch (minus gut content)	2.0
	White Perch (" " ")	1.8
	White Perch (" " ")	3.1
Lewis State Park, Ore.	Razor Clams (soft parts)	28
Illwaco, Washington	Chinook Salmon (dressed)	1.2
	Chinook Salmon (dressed)	2.2
	Starry Flounder (Minus gut content)	13
	Crab (flesh)	15
Long Beach, Wash.	Razor Clams (soft parts)	27
Willapa Bay, Wash.	Oysters (soft parts)	46
	Steamer Clams (soft parts)	1.3
Kalaloch, Wash.	Razor Clams (soft parts)	7.8
	Smelt (minus gut content)	14
Sequim, Wash.	Mixed Clams (soft parts)	0.74
	Oysters (soft parts)	4.2

* Data supplied by Radioecology Operation.

APPENDIX C-1

AVERAGE RADIOACTIVE PARTICLES CONCENTRATIONS*
AT SELECTED PACIFIC NORTHWEST LOCATIONS - 1959

Units of Particles per m³

Date	Rich- land	Proj- ect (2W)	Spokane	Boise	Klamath Falls	Lewis- ton	Walla Walla	Yakima
1-8	0.43	0.26	0.26	0.48	0.42	0.49	0.35	0.66
1-15	0.19	0.26	0.21	0.60	0.34	0.41	0.34	0.095
1-22	0.60	0.45	0.48	0.47	0.78	0.54	0.59	0.17
1-29	0.30	0.27	0.25	0.46	0.40	0.23	0.33	0.14
2-5	0.22	0.24	0.39	0.54	0.50	0.50	0.29	0.23
2-12	0.16	0.16	0.34	0.50	0.28	0.40	0.36	0.13
2-19	0.39	0.29	0.54	0.63	0.23	0.45	0.36	0.24
2-26	0.35	0.22	0.44	0.56	0.46	0.24	0.38	0.18
3-5	0.41	0.35	0.49	0.64	0.80	0.52	0.50	0.40
3-12	0.26	0.33	0.57	0.51	0.64	0.28	0.51	0.34
3-19	0.31	0.27	0.39	0.53	0.57	0.42	0.44	0.24
3-26	0.25	0.34	0.41	0.42	0.61	0.32	0.17	0.33
4-2	0.090	0.10	0.16	0.23	0.19	0.19	0.13	0.17
4-9	0.49	0.52	0.40	0.72	0.37	0.60	0.68	0.73
4-16	0.27	0.12	0.13	0.45	0.33	0.42	0.33	0.21
4-23	0.46	0.53	0.44	0.54	0.79	0.47	0.41	0.50
4-30	0.52	0.38	0.47	0.76	0.34	0.78	0.68	0.59
5-7	0.31	0.34	0.28	0.28	0.15	0.29	0.28	0.30
5-14	0.42	0.34	0.53	0.48	0.24	0.48	0.46	0.36
5-21	0.11	0.088	0.055	0.20	0.30	0.10	0.086	0.17
5-28	0.20	0.17	0.22	0.26	0.22	0.15	0.17	0.34
6-4	0.17	0.14	0.18	0.31	0.20	0.24	0.24	0.17
6-11	0.035	0.061	0.073	0.18	0.065	0.048	0.029	0.072
6-18	0.035	0.068	0.10	0.061	0.056	0.090	0.090	0.049
6-25	0.048	0.047	0.073	0.029	0.029	0.056	0.069	0.091
7-2	0.035	0.030	0.038	0.048	0.035	0.036	0.052	0.031
7-9	0.024	0.032	0.028	0.12	0.057	0.034	0.033	0.024
7-16	0.030	0.064	0.099	0.034	0.018	0.031	0.057	0.042

*Based on a nominal flow rate of 2.5 CFM

APPENDIX C-1 (CONTINUED)

<u>Date</u>	<u>Rich-land</u>	<u>Project (2W)</u>	<u>Spokane</u>	<u>Boise</u>	<u>Klamath Falls</u>	<u>Lewis-ton</u>	<u>Walla Walla</u>	<u>Yakima</u>
7-23	0.031	0.045	0.066	0.045	0.031	0.038	0.068	0.032
7-30	0.022	0.030	0.041	0.036	0.034	0.024	0.021	0.045
8-6	0.022	0.043	0.038	0.043	0.028	0.042	0.035	0.028
8-13	0.011	0.043	0.032	0.019	0.028	0.015	0.030	0.030
8-20	0.0097	0.012	0.011	0.020	0.0084	0.011	0.013	0.0097
8-27	0.0098	0.016	0.013	0.0098	0.0056	0.012	0.0097	0.011
9-3	0.0042	0.013	0.0099	0.0053	0.0084	0.0070	0.0083	0.0061
9-10	0.012	0.013	0.0056	0.019	0.0042	0.011	0.0073	0.0098
9-17	0.0085	0.0056	0.0042	0.0098	0.011	0.0052	0.0098	0.0097
9-24	0.0070	0.0042	0.0056	0.013	-0.0014	0.0028	0.0084	0.0017
10-1	0.0098	0.0042	0.0014	0.0085	0.0042	0.0014	0.0028	0.0028
10-8	0.0033	0.011	0.0070	0.0040	0.0045	0.0042	0.0098	0.0097
10-15	0.0036	0.011	0.0070	0.0088	0.0084	0.0087	0.0085	0.0085
10-22	0.0042	0.066	0.0028	0.015	0.0033	0.0054	0.0056	0.0048
10-29	0.0028	0.0042	0.0070	0.0098	0.0037	0.0014	0.0056	0.0056
11-5	0.0014	0.0042	0.0014	0.0028	0.0042	0.0041	-0.0014	0.0028
11-12	0.0056	0.0099	0.0055	0.012	0.0056	-0.0014	0.0056	0.0014
11-19	0.0028	0.0070	0.0014	0.0026	<0.0014	<0.0014	0.0042	0.0014
11-26	<0.0026	0.0033	0.0016	<0.0015	<0.0014	<0.0013	<0.0014	0.0014
12-3	0.0058	0.014	0.0014	0.0077	0.0085	0.0071	<0.0014	0.0062
12-10	0.0014	0.0097	<0.0014	0.0046	0.0014	0.0027	0.0056	0.0033
12-17	0.0028	0.0084	<0.0014	0.0026	0.0014	0.0057	0.0014	0.0070
12-24	<0.0014	0.011	<0.0016	0.0015	0.0014	0.0028	<0.0014	0.0084
12-31	<0.0014	<0.0014	<0.0014	<0.0014	0.0042	<0.0013	0.0014	<0.0014

APPENDIX C-2

AVERAGE BETA ACTIVITY ON FILTERS
FROM SELECTED NORTHWEST U. S. SAMPLING STATIONS - 1959

Beta Disintegrations per Second per cc of Filtered Air $\times 10^{-8}$

Date	Richland Wash.	Spokane Wash.	Boise Idaho	Hanford Project (200-W)	Klamath Falls Oreg.	Lewiston Idaho	Walla Walla Wash.	Yakima Wash.
1/8	1.3	15	18	12	17	19	14	18
1/15	8.2	11	36	12	13	21	12	7.3
1/22	17	13	21	14	25	22	16	19
1/29	12	8.4	17	7.5	14	11	12	6.5
2/5	9.3	9.6	18	6.2	13	15	8.5	8.1
2/12	6.9	8.3	14	4.4	9.5	12	7.8	6.5
2/19	12	12	20	7.7	7.8	15	11	8.6
2/26	11	11	24	7.5	24	15	12	10
3/5	15	18	41	16	35	24	23	19
3/12	10	11	20	7.8	26	14	12	9.8
3/19	10	12	26	6.3	22	13	13	7.9
3/26	13	18	26	9.5	15	15	13	10
4/2	5.0	5.5	12	3.3	7.5	9.6	5.6	5.0
4/9	17	12	32	12	17	22	18	15
4/16	18	16	28	12	26	21	16	19
4/23	20	18	23	14	19	25	15	20
4/30	17	20	24	12	20	36	24	14
5/7	16	15	23	15	12	17	13	13
5/14	20	19	31	13	19	25	18	16
5/21	8.0	5.3	15	5.4	19	10	7.8	6.3
5/28	11	12	18	8.2	14	12	8.5	12
6/4	14	11	24	9.2	18	18	13	11
6/11	5.8	7.3	13	4.7	7.2	9.6	5.4	5.5
6/18	8.9	8.9	6.3	7.0	7.2	12	6.8	7.5
6/25	7.5	7.3	6.4	5.2	4.9	11	7.8	8.3
7/2	4.9	4.9	8.5	3.6	5.6	7.4	5.1	4.5

APPENDIX C-2 (CONTINUED)

Date	Richland Wash.	Spokane Wash.	Boise Idaho	Hanford Project (200-W)	Klamath Falls Oreg.	Lewiston Idaho	Walla Walla Wash.	Yakima Wash.
7/9	2.6	3.0	8.1	1.9	6.0	4.2	3.5	3.3
7/16	4.6	5.0	6.5	3.2	2.9	6.3	4.4	3.6
7/23	4.6	4.5	5.1	3.1	3.2	6.3	4.4	3.8
7/30	3.1	2.9	3.5	2.3	4.7	4.2	3.0	2.8
8/6	3.1	3.1	4.4	2.4	3.1	5.5	3.8	2.6
8/13	2.6	2.5	3.1	2.0	2.4	3.1	3.0	2.4
8/20	1.4	1.4	2.8	0.89	1.7	2.1	1.3	1.6
8/27	1.5	1.1	2.0	1.1	1.3	2.1	1.2	1.3
9/3	1.1	1.3	1.7	0.82	0.92	1.8	1.4	1.1
9/10	1.1	0.69	1.6	0.60	0.65	1.1	0.97	0.83
9/17	1.4	1.1	1.2	0.90	0.97	1.8	1.7	1.1
9/24	0.88	0.78	0.74	0.49	1.0	2.0	2.4	0.36
10/1	0.66	0.47	0.75	0.73	0.69	0.56	0.81	0.64
10/8	1.9	1.1	0.82	1.2	1.1	1.6	1.4	0.93
10/15	1.1	0.72	0.84	1.1	0.83	1.2	1.3	0.76
10/22	1.3	0.91	1.5	1.4	1.0	1.1	1.6	1.1
10/29	1.1	0.64	0.99	1.1	0.55	1.2	1.2	1.1
11/5	1.4	0.93	1.2	1.4	0.69	1.2	1.2	1.1
11/12	2.1	0.92	1.0	2.3	0.63	1.7	1.7	2.0
11/19	1.2	0.70	1.0	1.5	1.0	1.4	1.3	1.1
11/26	0.39	0.54	0.66	1.2	0.39	0.33	0.51	0.33
12/3	0.97	0.49	1.1	2.2	1.2	1.2	1.1	0.97
12/10	1.5	0.43	1.5	1.6	0.80	1.1	0.82	1.6
12/17	0.89	0.48	0.79	0.88	0.40	0.75	0.60	0.63
12/24	2.4	0.31	1.2	1.6	0.53	0.74	0.78	0.65
12/31	1.6	0.46	0.90	1.2	0.87	0.71	0.86	0.72

APPENDIX C-3

ATMOSPHERIC CONCENTRATIONS OF I¹³¹

AT PERIMETER COMMUNITIES - 1959

$10^{-14} \mu\text{c I}^{131}/\text{cc}$

Date	Richland, Wash.	North Richland, Wash.	Date	Benton, Wash.	Pasco, Wash.
1/5	78	20	1/6	61	43
1/12	29	8.9	1/13	61	
1/19	4.8	20	1/15		44
1/26	21	20	1/19	24	
2/2	11	15	1/20		
2/9	9.8	13	1/27	33	
2/16	17	13	2/3	74	
2/23	23	8.8	2/10	100	
3/2	12	7.1	2/17	28	
3/9	6.3	13	2/24	32	
3/16	4.6	18	2/25	22	
3/23	3.5	7.7	3/3	45	
3/30	3.0	7.7	3/4		
4/6	6.9	4.9	3/10	59	
4/13	9.0	5.0	3/17	37	
4/20	6.3		3/24	31	
4/27	3.2		3/31	17	
5/4	1.6	6.4	4/7	15	
5/11	3.6	7.5	4/14	14	
5/18	2.5	8.7	4/21	12	
5/26	4.6	9.4	4/28	3.0	
6/1	0.3	4.6	4/29	5.2	
6/8	2.1	3.8	5/5	7.3	
6/15	1.5	3.3	5/12	8.5	
6/22	2.7	1.7	5/15	5.0	
6/29	1.0	4.7	5/19	5.6	
7/6	1.1	1.1	5/26	19	
7/13	3.0	0.8	6/2	6.3	
		1.7	6/9	4.9	

APPENDIX C-3 (CONTINUED)

<u>Date</u>	<u>Richland, Wash.</u>	<u>North Richland, Wash.</u>	<u>Date</u>	<u>Benton, Wash.</u>	<u>Pasco, Wash.</u>
7/20	22	1.8	6/16	4.6	25
7/27	3.5	1.9	6/23	4.1	2.5
8/3	2.1	1.4	6/30	1.8	2.8
8/10	3.5	5.1	7/7	3.2	0.7
8/17	1.8	2.9	7/14	11	0.6
8/24	5.7	3.1	7/21	2.2	3.7
8/31	8.1		7/28	1.4	2.7
9/8	1.3	1.7	8/4		1.6
9/14	3.8	3.0	8/6	4.4	
9/21	2.6	2.7	8/11	1.6	6.6
9/28	3.3	1.6	8/18	1.8	4.7
10/5	7.8	0.3	8/25	2.8	1.7
10/12	7.0	4.8	9/1	3.8	1.8
10/19	2.5	3.2	9/9	1.3	3.4
10/26	4.6	2.5	9/15	4.3	3.4
11/2	4.7	3.0	9/22	1.8	3.1
11/9	1.4	18	9/29	1.5	2.6
11/16	3.3	2.1	10/6	3.0	1.0
11/23	1.5	1.2	10/13	3.9	1.6
11/30	7.9	4.0	10/20	2.3	1.5
12/7	6.7	5.7	10/27	1.7	1.0
12/14	2.6	1.6	11/3	3.3	2.1
12/21	7.6	9.9	11/10	12	1.8
			11/17	1.5	3.4
			11/23		3.1
			11/24	3.1	
			11/30		10
			12/1	2.3	
			12/7		3.5
			12/8	11	
			12/14		1.6
			12/15	3.1	
			12/21		3.5
			12/22	21	

APPENDIX C-4

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES
BENTON CITY, WASHINGTON AND VICINITY, ZONE I - 1959
REPORTED IN UNITS OF 10^{-6} $\mu\text{c/g}$ VEGETATION

Sample Reporting Limits*	Ba-La 140	Zr-Nb 95	Ru 103+ Ru	Ce 144 Ce	Sample Date	Ba-La 140	Zr-Nb 95	Ru 103+ Ru	Ce 144 Ce
12/30/58	2.0	2.0	2.0	1.5	5.0	2.0	2.0	2.0	1.5
1/6/59	8.4	74	8.7	2.3	78	6/30	-	7.3	-
1/13	8.1	97	10	-	83	7/7	-	5.5	-
1/20	6.4	92	11	1.7	100	7/14	-	4.9	-
1/27	6.3	99	14	-	99	7/21	-	4.7	-
2/3	3.7	140	16	-	110	7/28	-	4.3	-
2/10	3.0	62	10	2.5	65	8/11	-	-	-
2/17	3.1	120	14	2.2	87	8/18	-	-	-
2/24	2.4	83	14	-	76	8/25	-	2.2	-
3/3	-	87	13	-	67	9/1	-	-	12
3/10	-	86	14	-	69	9/9	-	-	6.6
3/17	-	99	16	-	73	9/15	-	-	-
3/24	-	91	16	-	59	9/21	-	-	6.2
3/31	-	120	21	-	79	9/29	-	-	-
4/7	-	55	10	-	54	10/6	-	-	5.9
4/14	-	51	9.2	-	41	10/13	-	-	6.0
4/21	-	26	5.1	-	19	10/20	-	-	6.8
4/28	-	24	5.2	-	20	10/27	-	-	-
5/3	-	16	3.6	-	10	11/2	-	-	6.4
5/10	-	17	4.0	-	12	11/10	-	-	1.5
5/17	-	14	2.7	-	12	11/17	-	2.2	13
5/24	-	8.9	2.2	-	11/24	-	-	4.8	-
5/31	-	6.3	-	-	12/1	-	-	3.6	-
6/7	-	4.4	-	-	12/8	-	-	2.1	17
6/14	-	5.6	-	-	12/15	-	-	2.5	-
6/21	-	5.1	-	-	12/22	-	-	5.5	9.8
6/28	-	7.0	-	-	12/29	-	-	3.2	-

- 88 -

HW-64371

* Analytical results below reporting limits are indicated by a dash.

APPENDIX C-5

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES

RICHLAND, WASHINGTON AND VICINITY, ZONE K - 1959

REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

Sample Date	Ba-La ¹⁴⁰	Zr-Nb ⁹⁵	Ru ¹⁰³⁺ Ru ¹⁰⁶	I ¹³¹	Ce ¹⁴¹⁺ Ce ¹⁴⁴	Sample Date	Ba-La ¹⁴⁰	Zr-Nb ⁹⁵	Ru ¹⁰³⁺ Ru ¹⁰⁶	I ¹³¹	Ce ¹⁴¹⁺ Ce ¹⁴⁴
Reporting Limits*	2.0	2.0	2.0	1.5	5.0	2.0	2.0	2.0	2.0	1.5	5.0
12/29/58	8.1	71	9.3	-	81	7/6	-	2.8	-	-	-
1/5/59	5.0	51	10	-	58	7/13	-	4.2	-	-	5.8
1/12	7.6	66	9.1	-	80	7/20	-	4.6	-	-	-
1/19	6.2	100	14	-	92	7/27	-	2.1	-	-	-
1/26	7.3	97	21	-	110	8/3	-	2.4	-	-	-
2/2	6.9	72	12	2.8	75	8/10	-	-	-	-	-
2/9	3.7	110	16	3.2	110	8/17	-	-	-	-	-
2/16	2.2	110	22	2.3	94	8/24	-	2.6	-	-	6.4
2/25	8.5	220	42	-	220	8/31	-	-	-	-	-
3/2	3.8	150	39	-	140	9/8	-	-	-	-	-
3/9	2.2	190	26	-	88	9/14	-	-	-	-	-
3/16	-	120	24	-	140	9/21	-	-	-	-	6.8
3/23	-	88	15	-	57	9/28	-	-	-	-	-
3/30	-	40	8.0	-	31	10/5	-	-	-	-	6.8
4/6	-	44	11	-	36	10/12	-	-	-	-	5.3
4/13	-	41	7.3	-	34	10/19	-	-	-	-	-
4/20	-	28	6.8	-	23	10/26	-	-	-	-	10
4/27	-	20	5.6	-	13	11/2	-	-	-	-	-
5/4	-	26	5.8	-	16	11/9	-	-	-	-	-
5/13	-	3.8	12	2.0	22	11/16	-	2.8	-	3.0	1.9
5/19	-	24	5.0	-	13	11/25	-	4.1	-	-	9.7
5/25	-	8.7	-	-	-	12/2	-	2.3	-	-	9.9
6/1	-	5.3	-	-	-	12/9	-	-	-	2.8	7.0
6/8	-	7.7	-	-	-	12/16	-	2.7	-	11	-
6/15	-	9.5	-	-	-	12/23	-	2.4	6.8	9.6	-
6/23	-	9.6	-	-	-	12/30	-	4.7	10	-	-
6/29	-	4.2	-	-	-	-	-	-	-	-	-

* Analytical results below reporting limits are indicated by a dash.

APPENDIX C-6

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES

KENNEWICK, WASHINGTON AND VICINITY, ZONE L - 1959

REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

Reporting Limits*	Ba-La	140	Zr-Nb	95	Ru	$^{103+}$	Ce	$^{141+}$	Sample	Ru	$^{103+}$	Ru	106	I	131	Ce	$^{141+}$
	Date				Ru	106	Ce	144	Date		Ru	106	Ru		Ce	144	
2.0	2.0			2.0				5.0		2.0				2.0		5.0	
1/7	5.7	63		11				-	69	7/15				-		-	
1/14	5.9	76		9.8				-	92	7/22				-		-	
1/21	6.9	100		16				2.4	110	7/29				-		-	
1/28	5.6	110		16				-	110	8/5				-		-	
2/4	6.5	130		26				3.4	150	8/12				-		-	
2/11	4.2	130		22				1.6	170	8.19				-		-	
2/18	-	37		21				-	140	8/26				-		-	
2/24	2.8	100		21				-	100	9/2				-	10		
3/3	-	98		20				-	90	9/10				-		-	
3/11	2.0	100		22				-	94	9/16				-		-	
3/23	-	160		33				-	120	9/23				-	7.2		
3/31	-	100		22				-	88	9/30				-	2.3		
4/1	-	52		8.7				-	42	10/7				-	-	-	
4/8	2.0	44		9.5				-	38	10/14				-	-	-	
4/15	-	11		2.8				-	12	10/21				-	-	-	
4/22	-	14		3.1				-	11	10/28				-	-	-	
4/29	-	6.0		-				-	-	11/4				-	-	-	
5/6	-	6.9		2.7				-	7.0	11/11				-	3.1		
5/13	-	6.6		2.5				-	-	11/19				-	-	-	
5/20	-	14		3.9				-	8.9	11/24				-	2.4		
5/27	-	11		-				-	7.5	12/1				-	-	-	
6/3	-	6.0		-				-	5.0	12/8				-	2.2		
6/10	-	14		3.7				-	11	12/15				-	4.3		
6/17	-	6.4		-				-	5.2	12/22				-	6.8		
6/24	-	2.1		-				-	-	12/29				-	7.6		
7/1	-	3.4		-				-	-	-				-	3.4	1.8	
7/8	-	3.3		-				-	-	-				-	-	9.9	

* Analytical results below reporting limits are indicated by a (-).

APPENDIX C-7

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES
PASCO TO ELTOPIA, WASHINGTON, ZONE M - 1959
REPORTED IN UNITS OF 10^{-6} $\mu\text{c/g}$ VEGETATION

Sample Date	Ba-La ¹⁴⁰	Zr-Nb ⁹⁵	Ru ¹⁰³⁺ Ru ¹⁰⁶	Ru ¹³¹	Ce ¹⁴¹⁺ Ce ¹⁴⁴	Sample Date	Ba-La ¹⁴⁰	Zr-Nb ⁹⁵	Ru ¹⁰³⁺ Ru ¹⁰⁶	Ru ¹³¹	Ce ¹⁴¹⁺ Ce ¹⁴⁴
Reporting Limits*	2.0	2.0	2.0	1.5	5.0			2.0	2.0	1.5	5.0
1/7	3.8	66	9.7	-	61	7/8			2.6	-	-
1/14	6.7	94	12	-	92	7/15			2.6	-	-
1/21	5.3	110	11	1.9	91	7/22			-	-	-
1/28	7.8	130	19	-	140	7/29			-	-	-
2/4	6.3	140	27	-	170	8/5			2.5	-	-
2/11	6.2	190	38	2.9	190	8/12			-	-	-
2/18	6.0	150	28	-	160	8/19			-	-	-
2/24	2.5	91	13	-	70	8/26			-	-	-
3/3	-	120	27	-	110	9/2			2.1	2.1	12
3/11	2.0	130	23	-	110	9/10			2.3	2.5	8.9
3/18	-	120	21	-	89	9/16			-	-	-
3/25	-	93	17	-	67	9/23			-	-	5.9
4/1	-	56	8.6	-	53	9/30			-	-	-
4/8	-	42	6.2	-	37	10/7			-	-	-
4/15	-	33	6.1	-	30	10/14			-	-	-
4/22	-	49	8.6	-	41	10/21			-	-	-
4/29	-	18	4.0	-	16	10/28			2.0	-	-
5/6	-	25	6.7	-	16	11/4			-	-	-
5/13	-	16	4.0	-	11	11/11			-	-	-
5/20	-	17	3.3	-	14	11/19			-	3.7	3.3
5/27	-	8.4	-	-	6.6	11/24			2.4	-	5.4
6/3	-	5.1	-	-	-	12/1			-	-	3.4
6/10	-	12	3.4	-	9.0	12/8			4.6	-	6.3
6/17	-	6.3	-	-	5.2	12/15			2.3	-	13
6/24	-	2.4	-	-	-	12/22			-	-	-
7/1	-	3.5	-	-	-	12/29			4.4	-	-

* Analytical results below reporting limits are indicated by a (-).

HW-64371

APPENDIX C-8

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES
MESA, WASHINGTON AND VICINITY, ZONE N - 1959
REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

Sample Date	Ba-La 140	Zr-Nb 95	Ru 103+ Ru 106	I 131	Ce 141+ Ce 144	Sample Date	Ba-La 140	Zr-Nb 95	Ru 103+ Ru 106	Ru 131	Ce 141+ Ce 144
Reporting Limits*	2.0	2.0	2.0	1.5	5.0		2.0	2.0	2.0	1.5	5.0
1/7	4.6	65	12	-	69	7/8	-	-	3.9	-	-
1/14	5.2	78	11	-	87	7/15	-	-	6.1	-	-
1/21	3.7	75	7.8	-	58	7/22	-	-	3.4	-	-
1/28	5.9	110	22	-	130	7/29	-	-	-	-	-
2/4	7.8	140	28	-	170	8/5	-	-	-	-	-
2/11	7.8	170	31	-	110	8/12	-	-	-	-	-
2/18	4.9	150	32	-	170	8/19	-	-	-	-	-
2/24	2.1	100	20	-	110	8/26	-	-	-	-	-
3/3	4.7	140	28	-	130	9/2	-	-	2.5	-	-
3/11	-	140	28	-	130	9/10	-	-	5.0	-	-
3/18	2.2	160	29	-	100	9/16	-	-	-	-	-
3/25	-	150	20	-	95	9/23	-	-	-	-	-
4/1	-	38	6.8	-	33	9/30	-	-	-	-	-
4/8	-	28	4.2	-	18	10/7	-	-	-	-	-
4/15	-	27	4.3	-	22	10/14	-	-	-	-	-
4/22	-	14	3.5	-	15	10/21	-	-	2.6	-	-
4/29	-	15	3.3	-	11	10/28	-	-	-	-	-
5/6	-	16	3.4	-	11	11/4	-	-	2.4	-	-
5/13	-	23	3.2	-	15	11/11	-	-	-	-	-
5/20	-	13	2.9	-	12	11/19	-	-	-	-	-
5/27	-	4.1	-	-	11/24	2.0	-	-	-	-	-
6/3	-	4.8	-	-	12/1	-	-	-	-	-	-
6/10	-	9.8	2.1	-	7.9	12/8	-	-	-	-	-
6/17	-	3.4	-	-	-	12/15	-	-	2.1	-	-
6/24	-	5.1	-	-	-	12/22	-	-	2.4	-	-
7/1	-	5.5	-	-	-	12/29	-	-	5.4	-	-

* Analytical results below reporting limits are indicated by a (-).

APPENDIX C-9

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES

WAHLUKE SLOPE EAST, ZONE O - 1959
REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

Sample Date	Ba-La 140	Zr-Nb 95 Ru 103+ Ru 106 Ru	Ce 141+ Ce 144 Ce	Sample Date	Ba-La 140	Zr-Nb 95 Ru 103+ Ru 106 Ru	131 131	Ce 141+ Ce 144 Ce
Reporting Limits*	2.0	2.0	2.0	1.5	5.0	2.0	2.0	1.5
1/5	9.1	150	16	-	110	7/6	-	-
1/12	7.8	110	13	-	110	7/13	4.9	-
1/19	8.5	130	17	-	110	7/20	5.5	-
1/26	9.6	220	20	-	170	7/27	3.9	-
2/2	4.1	100	12	-	87	8/3	3.0	-
2/9	3.4	99	12	-	85	8/10	2.7	-
2/16	3.0	170	26	-	130	8/17	-	-
2/23	4.2	120	18	-	100	8/24	3.2	-
3/2	3.5	100	16	-	87	8/31	2.3	-
3/10	-	82	14	-	69	9/8	3.2	-
3/16	-	76	15	-	62	9/14	3.0	-
3/23	-	73	14	-	58	9/21	2.7	-
3/30	-	55	4.9	-	36	9/28	-	-
4/6	-	35	6.6	-	30	10/5	-	-
4/13	-	26	3.4	-	20	10/12	-	-
4/20	-	29	6.9	-	21	10/19	-	-
4/27	-	29	6.6	-	19	10/26	-	-
4/30	-	23	4.8	-	17	11/2	2.4	-
5/4	-	15	3.6	1.6	8.3	11/9	2.3	-
5/11	-	12	3.0	-	6.2	11/16	6.3	-
5/18	-	17	2.9	-	9.9	11/23	-	-
5/25	-	8.2	-	-	5.2	11/30	-	-
6/1	-	8.6	2.1	-	6.3	12/7	11	-
6/10	-	13	2.9	-	10	12/14	5.2	-
6/15	-	6.6	2.2	-	-	12/21	2.1	-
6/22	-	6.2	-	-	-	12/28	8.5	-
6/29	-	5.5	-	-	-	-	2.9	-

* Analytical results below reporting limits are indicated by a (-).

APPENDIX C-10

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES

WAHLUKE SLOPE WEST, ZONE P - 1959

REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

Reporting Limits*	Sample Date	Ba-La 140	Zr-Nb 95	Ru 103+ Ru 106	Ce 141+ Ce 144	Sample Date	Ba-La 140	Zr-Nb 95	Ru 103+ Ru 106	Ce 141+ Ce 144
1/5/59	5. 9	93	2. 0	-	1. 9	82	6/29	-	7. 7	-
1/12	7. 2	140	12	2. 7	120	7/6	-	4. 6	4. 6	-
1/19	5. 0	130	11	-	110	7/13	-	4. 6	4. 6	-
1/26	7. 5	210	11	-	140	7/20	-	4. 1	4. 1	-
2/2	3. 9	120	10	-	85	7/27	-	4. 3	4. 3	-
2/9	2. 5	100	12	1. 8	83	8/3	-	3. 6	3. 6	-
2/16	2. 0	110	12	1. 8	71	8/10	-	2. 2	2. 2	-
2/23	3. 1	110	23	-	99	8/17	2. 0	2. 1	-	-
3/2	-	96	16	-	77	8/24	-	3. 4	2. 4	-
3/10	-	74	83	-	47	8/31	-	2. 3	-	-
3/16	-	68	11	-	47	9/8	-	-	-	-
3/23	-	58	7. 6	-	32	9/14	-	-	-	-
3/30	-	42	5. 9	-	32	9/21	-	-	-	-
4/6	-	38	4. 9	-	30	9/28	-	-	-	-
4/13	-	20	3. 7	-	16	10/5	-	-	-	-
4/20	-	27	6. 9	-	22	10/12	-	-	-	-
4/27	-	30	6. 8	-	16	10/19	-	-	-	-
4/30	-	15	3. 9	-	9. 3	10/26	-	-	-	-
5/4	-	22	4. 6	-	13	11/2	-	-	2. 8	-
5/11	-	12	2. 3	-	8. 3	11/9	-	-	3. 2	-
5/18	-	17	2. 5	-	8. 9	11/16	-	2. 9	3. 5	-
5/25	-	9. 0	2. 3	-	-	11/23	-	-	1. 5	-
6/2	-	8. 0	-	-	5. 8	12/14	-	-	7. 0	-
6/10	-	10	3. 2	-	8. 6	12/21	-	-	-	-
6/15	-	8. 3	2. 5	-	13	12/28	-	-	2. 9	-
6/22	-	6. 8	-	-	5. 0	-	-	-	-	-

* Analytical results below reporting limits are indicated by a (-).

APPENDIX C-11

CONCENTRATION OF SELECTED RADIONUCLIDES ON NATIVE GRASSES

HORSE HEAVEN HILLS, ZONE Q - 1959

REPORTED IN UNITS of $10^{-6} \mu\text{c/g}$ VEGETATION

Sample Reporting Limits*	Ba-La 140	Zr-Nb 95	Ru 103+ Ru 106	I 131	Ce 141+ Ce 144	Sample Date	Ba-La 140	Zr-Nb 95	Ru 103+ Ru 106	I 131	Ce 141+ Ce 144
12/22/58	2.0	2.0	2.0	1.5	5.0		2.0	2.0	2.0	1.5	5.0
12/30	17	97	13	-	100	7/9	-	3.4	-	-	-
1/15/59	8.7	61	9.8	-	58.	7/16	-	-	-	-	-
1/22	9.0	120	22	-	160	7/23	-	-	-	-	-
1/29	10	120	25	1.7	130	7/30	-	4.7	-	5.0	-
1/29	11	190	31	-	180	8/6	-	-	-	-	-
2/5	4.1	140	25	3.0	140	8/13	-	4.2	-	-	-
2/12	5.4	120	23	-	120	8/20	-	-	2.5	-	-
2/19	4.5	170	35	-	160	8/27	-	2.6	-	12	6.5
2/26	-	97	21	2.4	100	9/3	-	-	-	-	3.1
3/5	4.4	180	31	-	170	9/11	-	-	-	-	5.2
3/12	-	180	35	-	150	9/17	-	-	-	-	5.3
3/26	-	120	22	-	90	9/24	-	-	-	-	-
4/1	-	48	8.2	-	41	10/1	-	-	-	-	4.8
4/9	-	49	10	-	46	10/8	-	-	-	-	-
4/16	-	37	8.3	-	30	10/15	-	-	-	-	-
4/23	-	29	5.5	-	21	10/22	-	-	-	-	-
5/7	-	8.7	2.6	-	8.2	10/29	-	-	-	-	5.7
5/14	-	15	2.7	-	7.7	11/5	-	-	-	-	7.6
5/21	-	13	3.1	-	7.8	11/11	-	-	-	-	8.7
5/27	-	15	2.9	-	8.0	11/19	-	-	-	-	-
6/4	-	9.7	2.1	-	5.9	11/23	-	-	2.1	-	5.6
6/11	-	7.7	2.0	-	7.9	12/2	-	-	3.6	-	-
6/18	-	5.1	-	-	-	12/9	-	-	3.0	-	5.5
6/25	-	-	-	-	-	12/16	-	-	2.0	-	11
7/2	-	2.4	-	-	-	12/23	-	-	-	-	3.1

* Analytical results below reporting limits are indicated by a dash.

HW-64371

-95-

APPENDIX C-12

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES
WALLA WALLA, WASHINGTON AND VICINITY, ZONE R - 1959

REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

Sample Date	Ba-La ¹⁴⁰	Zr-Nb ⁹⁵	Ru ¹⁰³⁺ Ru ¹⁰⁶	I ¹³¹	Ce ¹⁴¹⁺ Ce ¹⁴⁴
Reporting Limits *	2.0	2.0	2.0	1.5	5.0
12/9/58	29	74	13	-	120
1/13/59	14	130	29	-	180
2/10	8.6	160	29	-	170
3/10	3.4	280	45	-	250
4/15	-	23	5.6	-	20
4/28	-	33	7.2	-	27
5/19	-	21	5.1	-	12
7/8	-	2.8	-	-	-
8/18	-	7.9	4.9	-	27
9/1	-	-	4.3	-	8.0
10/6	-	-	-	-	-
10/27	-	-	-	-	-

* Analytical results below reporting limits are indicated by a dash.

APPENDIX C-13

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES

LEWISTON, IDAHO AND VICINITY, ZONE S - 1959

REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

<u>Sample Date</u>	<u>Ba-La¹⁴⁰</u>	<u>Zr-Nb⁹⁵</u>	<u>Ru¹⁰³⁺ Ru¹⁰⁶</u>	<u>I¹³¹</u>	<u>Ce¹⁴¹⁺ Ce¹⁴⁴</u>
Reporting Limits*	2.0	2.0	2.0	1.5	5.0
1/13	31	240	55	-	330
2/10	5.5	120	27	-	140
3/10	2.4	250	45	-	210
4/15	-	27	4.5	-	19
4/28	-	21	3.8	-	13.7
5/19	-	12	2.4	-	9.9
7/7	-	7.8	-	-	5.3
8/18	-	4.4	-	-	7.8
9/1	-	-	4.6	-	7.9
10/6	-	-	-	-	7.7
10/27	-	-	-	-	-

* Analytical results below reporting limits are indicated by a dash.

APPENDIX C-14

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES
SPOKANE, WASHINGTON AND VICINITY, ZONE T - 1959

REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

<u>Sample Date</u>	<u>Ba-La¹⁴⁰</u>	<u>Zr-Nb⁹⁵</u>	<u>Ru¹⁰³⁺/ Ru¹⁰⁶</u>	<u>I¹³¹</u>	<u>Ce¹⁴¹⁺/ Ce¹⁴⁴</u>
Reporting Limits*	2.0	2.0	2.0	1.5	5.0
8/5/58	11	20	4.8	-	28
8/25/58	2.8	11	3.5	-	14
1/14/59	18	210	40	-	280
2/11	10	230	44	-	240
3/10	4.6	250	42	-	200
4/15	-	41	6.4	-	35
4/28	-	47	8.7	-	32
5/20	-	17	3.1	-	12
7/8	-	-	-	-	-
8/19	-	9.7	-	-	16
9/1	-	3.8	4.5	-	15
10/6	-	2.2	-	-	12
10/27	-	-	-	-	-

* Analytical results below reporting limits are indicated by a dash.

APPENDIX C-15

CONCENTRATION OF SELECTED RADIONUCLIDES ON NATIVE GRASSES
RITZVILLE, WASHINGTON AND VICINITY, ZONE U - 1959

REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

<u>Sample Date</u>	<u>Ba-La¹⁴⁰</u>	<u>Zr-Nb⁹⁵</u>	<u>Ru103+ Ru¹⁰⁶</u>	<u>I¹³¹</u>	<u>Ce¹⁴¹⁺ Ce¹⁴⁴</u>
Reporting Limits *	2.0	2.0	2.0	1.5	5.0
1/14	12	130	18	-	160
2/12	5.1	200	33	-	170
3/11	3.4	230	37	-	200
4/15	-	32	4.8	-	27
4/29	-	11	2.5	-	8.8
5/20	-	10	2.2	-	8.8
7/8	-	6.7	-	-	-
8/19	-	3.1	-	-	-
9/2	-	-	-	-	6.3
10/7	-	-	-	-	6.6
10/28	-	-	-	-	-

* Analytical results below reporting limits are indicated by a dash.

APPENDIX C-16

CONCENTRATION OF SELECTED RADIONUCLIDES ON NATIVE GRASSES
TOPPENISH, WASHINGTON AND VICINITY, ZONE V - 1959
REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

<u>Sample Date</u>	<u>Ba-La¹⁴⁰</u>	<u>Zr-Nb⁹⁵</u>	<u>Ru¹⁰³⁺/ Ru¹⁰⁶</u>	<u>I¹³¹</u>	<u>Ce¹⁴¹⁺/ Ce¹⁴⁴</u>
Reporting Limits*	2.0	2.0	2.0	1.5	5.0
10/7/58	-	7.7	-	-	8.0
12/2/58	8.4	20	4.4	-	33
1/7/59	16	94	17	-	100
2/3	3.3	81	13	-	81
3/3	8.3	120	19	-	96
4/8	-	18	3.0	-	14
4/20	-	28	5.4	-	22
5/12	-	2.9	-	-	-
6/23	-	3.4	-	-	-
7/28	-	-	-	-	-
8/25	-	-	-	-	6.1
9/29	-	-	-	-	-
10/19	-	-	-	-	5.4
11/4	-	-	-	-	-
12/1	-	-	2.6	-	-

* Analytical results below reporting limits are indicated by a dash.

APPENDIX C-17

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES
GOLDENDALE, WASHINGTON AND VICINITY, ZONE W - 1959
REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

<u>Sample Date</u>	<u>Ba-La¹⁴⁰</u>	<u>Zr-Nb⁹⁵</u>	<u>Ru¹⁰³⁺/ Ru¹⁰⁶</u>	<u>I¹³¹</u>	<u>Ce¹⁴¹⁺/ Ce¹⁴⁴</u>
Reporting Limits*	2.0	2.0	2.0	1.5	5.0
10/7/58	2.0	4.0	-	-	5.9
12/2/58	32	81	15	-	120
1/7/59	12	82	14	-	97
2/3	18	340	56	-	420
3/3	-	320	62	-	310
4/8	2.2	65	10	-	64
4/20	-	44	10	-	39
5/12	-	21	3.9	-	15
6/23	-	3.3	-	-	-
7/28	-	2.3	-	-	-
8/25	-	3.0	4.5	-	-
9/29	-	-	-	-	-
10/19	-	-	2.7	-	-
11/4	-	-	-	-	6.8
12/1	-	-	-	-	5.4

* Analytical results below reporting limits are indicated by a dash.

APPENDIX C-18

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES
PORLAND TO BONNEVILLE, OREGON, ZONE X - 1959
REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

<u>Sample Date</u>	<u>Ba-La¹⁴⁰</u>	<u>Zr-Nb⁹⁵</u>	<u>Ru¹⁰³⁺ Ru¹⁰⁶</u>	<u>I¹³¹</u>	<u>Ce¹⁴¹⁺ Ce¹⁴⁴</u>
Reporting Limits*	2.0	2.0	2.0	1.5	5.0
10/7/58	2.0	9.8	-	-	11
12/2/58	27	54	13	-	94
1/8/59	15	100	24	-	150
2/3	28	460	85	-	590
3/3	9.8	540	82	-	500
4/9	-	130	22	-	130
4/20	-	29	6.6	-	24
5/13	-	17	2.7	-	12
6/24	-	7.1	-	-	5.1
7/29	-	2.2	-	-	-
8/26	-	3.9	-	-	-
9/29	-	5.4	2.7	-	22
10/19	-	3.8	-	-	27
11/4	-	-	-	-	-
12/1	-	2.3	2.0	-	9.9

* Analytical results below reporting limits are indicated by a dash.

APPENDIX C-19

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES
THE DALLES TO BOARDMAN, OREGON, ZONE Y - 1959
REPORTED IN UNITS OF $10^6 \mu\text{c/g}$ VEGETATION

Sample Date	Ba-La ¹⁴⁰	Zr-Nb ⁹⁵	Ru ¹⁰³⁺ Ru ¹⁰⁶	I ¹³¹	Ce ¹⁴¹⁺ Ce ¹⁴⁴
Reporting Limits*	2.0	2.0	2.0	1.5	.5.0
10/8/58	-	6.0	-	-	.8.3
12/3/58	18	61	7.6	-	75
1/8/59	3.7	59	9.7	-	58
2/4	4.7	130	17	-	120
3/4	4.8	170	31	-	140
4/9	-	35	7.1	-	30
4/20	-	37	8.1	-	26
5/13	-	17	3.9	-	11
6/24	-	4.5	-	-	-
7/29	-	-	-	-	-
8/26	-	2.2	-	-	6.2
9/30	-	-	-	-	6.2
10/20	-	-	-	-	10
11/4	-	3.7	-	-	21
12/2	-	-	4.0	-	-

*Analytical results below reporting limits are indicated by a dash.

APPENDIX C-20

CONCENTRATIONS OF SELECTED RADIONUCLIDES ON NATIVE GRASSES
 RIVERVIEW DISTRICT OF PASCO, WASHINGTON, ZONE Z - 1959
 REPORTED IN UNITS OF $10^{-6} \mu\text{c/g}$ VEGETATION

Sample	Ba-La	Zn ⁶⁵	Zr-Nb ⁹⁵	Cs ¹³⁷	Ru ¹⁰³⁺ Ru ¹⁰⁶	I ¹³¹	Cr ⁵¹	Ce ¹⁴¹⁺ Ce ¹⁴⁴
Reporting Limits*	2.0	1.5	2.0	1.0	2.0	1.5	5.0	5.0
12/31/58	9.5	-	58	3.4	8.7	-	-	66
1/7/59	6.0	-	50	4.5	10	-	-	66
1/14	9.9	-	61	6.4	11	-	-	87
1/21	9.5	4.1	110	9.1	19	-	-	130
1/28	4.1	5.6	74	4.7	15	-	-	78
2/4	5.1	-	39	7.9	26	3.4	-	150
2/11	6.1	1.5	110	7.1	21	-	-	110
2/18	5.1	2.9	140	14	26	-	-	150
2/24	5.1	4.8	120	7.8	22	-	-	120
3/3	6.4	6.1	160	9.2	31	-	-	130
3/11	3.0	4.5	200	9.0	39	-	-	160
3/18	-	2.2	160	9.1	30	-	-	130
3/25	14	9.1	200	14	40	-	-	150
4/1	-	2.3	49	3.8	9.2	-	-	41
4/9	-	1.8	47	4.5	10	-	-	40
4/15	-	1.7	14	1.9	2.2	-	-	9.1
4/22	-	3.0	11	1.3	3.0	-	-	-
4/29	-	1.8	4.1	-	10	2.2	-	5.0
5/6	-	1.5	7.2	-	-	-	-	-
5/13	-	-	7.1	-	-	-	-	-
5/20	-	-	8.3	-	-	-	-	-
5/27	-	-	8.5	-	-	-	-	-

* Analytical results below reporting limits are indicated by a dash.

APPENDIX C-20 (contd.)

Sample Date	Ba-La ¹⁴⁰	Zn ⁶⁵	Zr-Nb ⁹⁵	Cs ¹³⁷	Ru ¹⁰³⁺ Ru ¹⁰⁶	I ¹³¹	Cr ⁵¹	Ce ¹⁴¹⁺ Ce ¹⁴⁴
Reporting Limits*	2.0	1.5	2.0	1.0	2.0	1.5	5.0	5.0
6/3	-	2.6	8.1	-	-	-	-	-
6/10	-	3.0	5.2	-	-	-	-	-
6/17	-	-	5.2	-	-	-	-	-
6/24	-	-	3.0	-	-	-	-	-
7/1	-	2.5	-	-	-	-	-	-
7/8	-	2.6	2.3	-	-	-	-	-
7/15	-	1.5	-	-	-	-	-	-
7/22	-	-	-	-	-	-	-	-
7/29	-	2.6	4.0	-	-	-	-	-
8/5	-	-	2.1	-	-	-	-	-
8/12	-	-	-	-	-	-	-	-
8/19	-	-	-	-	-	-	-	-
8/26	-	-	-	-	-	-	-	-
9/2	-	1.9	12	1.4	-	2.1	-	-
9/10	-	-	2.1	-	-	2.1	-	6.0
9/16	-	-	2.9	-	-	-	-	13
9/23	-	3.6	-	1.5	2.9	-	-	-
9/30	-	-	6.2	1.1	3.1	-	-	25
10/7	-	-	-	-	-	-	-	-
10/14	-	-	-	2.1	2.1	-	-	-
10/21	-	-	-	-	3.7	-	-	-
10/28	-	-	2.8	-	-	-	-	-
11/4	-	5.3	-	-	-	-	-	-
11/11	-	2.9	2.4	1.5	-	-	-	5.1
11/24	-	3.7	-	-	-	-	-	9.2
12/1	-	11.	-	-	-	-	-	-
12/8	-	3.1	1.9	4.4	-	-	-	10
12/15	-	3.4	-	-	-	-	-	18
12/22	-	2.0	2.0	-	3.8	-	-	-
12/29	-	-	-	-	-	-	-	15

* Analytical results below reporting limits are indicated by a dash.

APPENDIX C-21
QUANTITY OF I^{131} RELEASED
FROM THE SEPARATIONS AREAS' PROCESS STACKS - 1959

Day of Month	January	February	March	April	May	June	July	August	September	October	November	December
1	0.03	2.6	0.47	0.14	1.5	0.63	0.69	0.61	0.41	0.95	0.96	0.22
2	0.023	2.8	0.29	0.38	1.4	0.81	0.46	0.61	0.52	0.28	0.92	1.2
3	0.023	1.2	0.43	0.20	1.4	0.99	0.46	0.64	0.38	0.28	0.82	0.29
4	0.023	0.68	0.23	0.40	1.7	1.8	0.34	0.57	0.40	0.28	0.89	0.45
5	0.24	0.80	0.21	0.19	0.35	2.3	0.33	0.65	0.40	0.25	0.21	0.87
6	0.25	0.72	0.33	0.22	0.66	2.3	0.28	0.69	0.29	0.30	0.92	0.87
7	0.31	0.72	0.33	0.33	0.80	2.3	0.23	1.5	0.23	0.34	1.6	0.92
8	0.54	0.72	0.33	0.64	1.5	2.5	0.27	2.8	0.29	0.24	1.6	1.2
9	0.53	0.27	0.22	0.87	1.7	1.2	0.23	2.8	0.44	0.47	2.0	1.6
10	0.53	0.79	0.48	1.0	1.7	1.0	0.20	0.50	0.35	0.48	2.0	0.45
11	0.53	0.42	0.43	1.6	1.4	0.84	0.19	0.65	0.40	0.47	0.95	0.52
12	0.52	0.11	0.32	1.6	1.3	0.99	0.19	0.24	0.41	0.19	0.82	0.52
13	0.68	0.008	0.60	1.2	1.1	0.99	0.39	0.33	0.41	0.27	1.0	0.51
14	0.34	0.009	0.70	1.5	1.2	0.93	0.37	0.22	0.28	0.49	1.0	0.49
15	1.1	0.010	0.70	1.0	1.3	0.87	0.37	0.21	0.093	0.60	1.0	0.77
16	0.27	0.020	0.84	1.0	0.78	0.83	0.33	0.21	0.41	0.51	1.2	0.99
17	0.003	0.030	0.84	1.1	0.79	0.65	0.34	0.23	0.50	0.51	1.3	1.0
18	0.004	0.019	0.31	1.4	0.76	0.63	0.38	0.20	0.80	0.52	1.3	0.87
19	0.42	0.029	0.58	1.4	0.40	0.62	0.38	0.55	0.83	0.33	0.48	1.2
20	0.45	0.042	0.42	1.2	1.5	0.59	0.34	0.53	0.83	0.58	0.96	1.2
21	0.61	0.042	0.43	1.0	1.3	0.59	1.0	0.84	0.95	0.37	0.96	1.2
22	0.81	0.051	0.43	0.61	1.1	0.73	0.87	0.88	0.72	0.58	0.98	1.4
23	0.90	0.084	0.43	0.67	1.1	0.55	0.65	0.89	1.1	0.44	1.4	1.6
24	0.65	0.060	0.36	2.6	1.0	0.66	1.1	0.94	0.72	0.60	1.5	1.1
25	0.65	0.11	0.61	2.6	0.25	0.71	1.1	0.89	1.4	0.60	0.51	1.1
26	0.62	0.74	0.36	2.6	0.22	0.64	1.1	0.89	1.4	0.64	0.52	1.5
27	0.72	0.47	0.06	2.5	0.16	0.43	0.77	0.59	1.4	0.77	0.43	1.5
28	0.99	0.46	0.06	1.4	0.26	0.43	0.73	0.83	0.64	0.87	0.35	1.5
29	1.4	0.06	1.4	0.37	0.16	1.4	0.79	0.79	1.4	0.35	1.2	1.1
30	1.9	0.01	1.7	0.65	0.10	0.93	0.10	0.61	1.5	0.33	1.1	1.2
31	2.6	0.03	0.03	0.65	0.65	0.34	0.34	0.34	0.96	0.96		

APPENDIX C-22
CONCENTRATION OF I¹³¹ IN THYROID TISSUE
OF JACK RABBITS* - 1959
Units - $10^{-6} \mu\text{c I}^{131}/\text{g (wet)}$

Date	Number of Samples	Location **	Thyroid		Muscle Average
			Average	Maximum	
1/12	4	Zone J	1800	4100	6.5
1/12	4	E	1400	2600	9.1
1/13	2	O	960	1900	12
2/ 9	4	J	2700	4800	8.1
2/ 9	4	E	2000	3200	5.6
2/10	2	O	700	910	6.6
3/ 4	4	E	430	1000	9.7
3/ 5	4	O	880	1900	7.2
4/ 1	4	J	81	220	5.9
4/ 1	4	E	200	620	5.5
4/ 2	2	O	200	590	2.2
5/ 5	3	J	300	380	4.6
5/ 5	2	E	390	560	3.2
5/ 6	3	O	280	410	6.4
6/11	4	J	240	490	6.1
6/11	4	E	280	620	12
6/12	4	O	380	940	9.4
7/ 7	4	J	390	690	44
7/ 7	4	E	230	970	18
7/ 8	4	O	220	690	17
8/ 5	4	J	130	160	6.9
8/ 5	4	E	210	370	6.2
8/ 6	4	O	65	84	7.9
9/ 1	4	J	230	510	5.0
9/ 1	4	E	110	240	4.2
9/ 2	4	O	150	790	5.3
10/ 5	4	J	1400	2000	5.1
10/ 5	4	E	1800	4600	5.6
10/ 6	4	O	220	350	9.7
11/10	4	E	2200	3800	14
11/11	3	J	1300	2000	6.6
11/11	4	O	1300	4300	6.5
12/ 7	2	J	1800	2800	5.3
12/ 7	3	E	2600	6400	5.2
12/ 8	3	O	2000	2400	15

* Data provided by Radioecology Operation, W. C. Hansen

** Refer to Figure 12 in text

APPENDIX D-1

CONCENTRATIONS OF SEVERAL RADIONUCLIDES IN MILK PURCHASED
FROM PRODUCERS AT SELECTED LOCATIONS - 1959

$10^{-6} \mu\text{c/g}$

<u>Code</u>	<u>Date</u>	<u>K⁴⁰</u>	<u>Zn⁶⁵</u>	<u>Cs¹³⁷</u>	<u>Sr⁸⁹</u>	<u>Sr⁹⁰</u>
Reporting Limits*		0.3	0.05	0.05	0.004	0.002

Riverview Irrigation District

Farm 1

54	1/ 5	0.81	-	-	-	0.0021
57	2/ 4	1.1	-	-	-	-
59	2/18	0.98	-	-	0.019	-
62	3/11	1.4	-	0.072	0.042	-
65	3/23	0.93	-	0.062	0.036	0.0070
69	4/ 1	1.2	-	0.066	0.062	0.0065
84	4/22	0.71	-	0.048	Lost	Lost
91	4/28	1.3	-	0.058	-	-
117	5/25	0.98	-	0.037	0.021	0.0029
121	6/24	0.84	-	0.036	0.0066	0.0024

Farm 2

109	5/25	1.3	0.70	0.020	0.0094	0.0025
124	6/30	0.96	0.34	0.035	0.0080	0.0048
387	11/19	1.5	0.58	-	Analysis not Completed	

Ringold Farms

Farm 1

305	9/24	1.6	0.42	-	0.0060	0.0020
-----	------	-----	------	---	--------	--------

* Results less than the reporting limit are indicated by a (-).

APPENDIX D-1 (contd.)

<u>Code</u>	<u>Date</u>	<u>K⁴⁰</u>	<u>Zn⁶⁵</u>	<u>Cs¹³⁷</u>	<u>Sr⁸⁹</u>	<u>Sr⁹⁰</u>
<u>Local Purchase - Commercial Milk</u>						
<u>Brand A</u>						
55	1/14	0.76	-	0.028	0.0059	0.0038
56	2/ 3	0.75	-	0.022	0.0072	0.0028
58	2/17	0.74	-	0.028	0.0084	-
63	3/11	0.79	-	0.022	0.0040	0.0040
64	3/18	0.86	-	0.036	0.0080	0.0040
72	4/ 9	0.64	-	0.032	0.014	0.0040
74	4/16	1.1	-	0.040	Lost	Lost
82	4/22	0.81	-	0.036	0.027	-
114	5/25	1.1	-	0.036	Lost	Lost
327	9/29	1.2	-	0.024	-	0.0027
386	11/17	1.4	-	-	Analysis not Completed	
413	12/ 8	1.2	-	-	"	"
<u>Brand B</u>						
88	4/27	1.2	-	0.032	Lost	Lost
110	5/25	1.0	-	0.036	0.018	0.0042
329	9/30	1.0	-	0.038	-	0.0020
382	11/17	1.5	-	0.040	-	0.0035
417	12/ 8	1.3	-	-	Analysis not Completed	
<u>Brand C</u>						
85	4/22	1.6	-	0.032	-	-
115	5/25	0.88	-	0.045	0.0072	0.004
230	8/19	1.4	-	-	-	0.002
325	9/29	1.2	-	-	0.0042	0.0038
384	11/17	1.4	-	0.026	Analysis not Completed	
415	12/ 8	0.92	-	-	"	"

Results less than the reporting limit are indicated by a (-).

APPENDIX D-1 (contd.)

<u>Code</u>	<u>Date</u>	<u>K⁴⁰</u>	<u>Zn⁶⁵</u>	<u>Cs¹³⁷</u>	<u>Sr⁸⁹</u>	<u>Sr⁹⁰</u>
Reporting Limits*		0.3	0.05	0.02	0.004	0.002
Brand D						
90	4/27	1.4	-	0.043	-	-
116	5/25	0.96	-	0.029	0.0087	0.0050
331	9/30	1.4	-	-	0.0086	0.0036
419	12/ 8	0.93	-	-	Analysis not Completed	
Brand E						
89	4/27	1.2	-	0.033	0.014	-
111	5/25	1.0	-	0.036	0.029	0.0072
330	9/30	1.5	-	0.044	Analysis not Completed	
Brand F						
83	4/22	0.62	-	0.020	Lost	Lost
113	5/25	0.87	-	-	0.018	-
332	9/30	1.5	-	-	0.0080	0.0030
381	11/17	0.93	-	-	-	0.0040
418	12/ 8	1.0	-	-	Analysis not Completed	
Brand G						
101	5/19	1.2	-	0.025	Lost	Lost
112	5/25	1.2	-	0.058	0.019	0.0056
228	8/19	1.2	-	-	-	-
326	9/29	1.1	-	0.025	0.0086	0.0036
385	11/17	1.4	-	-	Analysis not Completed	
414	12/ 8	1.5	-	-	"	"

* Results less than the reporting limit are indicated by a (-).

APPENDIX D-1 (contd.)

<u>Code</u>	<u>Date</u>	<u>K⁴⁰</u>	<u>Zn⁶⁵</u>	<u>Cs¹³⁷</u>	<u>Sr⁸⁹</u>	<u>Sr⁹⁰</u>
Reporting Limits*		0.3	0.05	0.02	0.004	0.002

Local Purchase - Coastal Origin - Commercial Milk

Brand H

66	3/31	1.3	-	0.073	0.012	0.007
71	4/ 9	Lost	Lost	Lost	-	-
81	4/22	0.98	-	0.091	Lost	Lost
107	5/25	1.0	-	0.091	0.086	0.018
125	6/30	0.77	-	0.080	0.035	0.013
163	6/30	1.5	-	0.084	0.018	0.0082
229	8/19	1.3	-	-	0.0035	0.0020
324	9/29	1.4	-	0.072	0.0097	0.0117
383	11/17	1.3	-	-	Analysis not Completed	
416	12/ 8	0.9	-	0.057	"	"

*Results less than the reporting limit are indicated by a (-).

APPENDIX D-2

CONCENTRATIONS OF SEVERAL RADIONUCLIDES
IN LOCALLY PURCHASED MODIFIED MILK - 1959

$10^{-6} \mu\text{c/g}$

Code	Brand	Date	Sc ⁴⁶	K ⁴⁰	Cs ¹³⁷	I ¹³¹	Cr ⁵¹	Ce ¹⁴¹⁺¹⁴⁴
Reporting Limits *			0.1	0.3	0.02	0.1	0.5	0.5
<u>Condensed or Evaporated Milk</u>								
450	C	12/23/59	-	3.0	0.17	-	-	-
451	J	12/23/59	-	1.7	0.041	-	-	-
452	K	12/23/59	-	2.7	0.081	-	-	-
453	L	12/23/59	-	2.8	0.13	-	-	4.6
<u>Dry or Powdered Milk</u>								
333	K	9/30/59	-	14	0.64	-	-	-
380	K	11/17/59	-	14	0.30	-	-	-
391	K	11/23/59	-	15	0.57	0.1	-	-
412	K	12/ 8/59	-	14	0.50	-	-	-
389	C	11/23/59	0.12	14	0.48	0.1	-	-
411	C	12/ 8/59	-	14	0.16	-	-	-
393	H	11/23/59	0.11	14	0.46	-	0.62	-
421	H	12/ 8/59	-	14	0.40	-	-	1.9
392	M	11/23/59	0.12	16	0.52	0.1	-	-
395	N	11/23/59	0.14	13	0.37	-	-	-

* Results less than the reporting limit are indicated by a (-).
 No results for Zr-Nb⁹⁵ or Ru¹⁰³⁺¹⁰⁶ were found above their respective reporting limits of 0.1 and $0.5 \times 10^{-6} \mu\text{c/g}$. The strontium analyses are not completed.

APPENDIX D-3

CONCENTRATIONS OF SEVERAL RADIONUCLIDES IN CEREAL CROPS PURCHASED FROM GROWERS

Code	Crop	Reporting Limits	Location Code	Date	AT SELECTED LOCATIONS - 1959 IN UNITS OF $10^{-6} \mu\text{c}/\text{g}$									
					Sc 46	K 40	Zn 65	Zr 95-Nb 95	Cs 137	Ru 103+ Ru 106	Ru 131	Cr 51	Ce 141+ Ce 144	Sr 89
Riverview Irrigation District														
146	Wheat	Hs	7/16	NA	7.3	1.9	1.3	-	1.1	-	0.98	1.8	0.083	0.036
169	"	Hs	7/22	NA	4.7	1.8	0.63	-	0.53	-	0.58	0.036	0.012	
170	Corn	Hs	7/22	NA	2.8	1.7	-	-	-	-	4.4	-	-	
194	"	Hs	8/10	NA	2.7	1.2	-	-	-	-	1.1	-	-	
206	"	Hs	8/19	NA	-	-	-	-	-	-	-	-	-	
238	"	Pr	8/27	NA	3.1	2.2	-	-	-	-	-	-	-	
240	"	Pr	9/1	NA	2.7	1.6	-	-	-	-	-	-	-	
244	"	Pr	9/9	NA	2.1	1.6	-	-	-	-	-	-	-	
247	Wheat	Cd	9/9	NA	5.5	1.4	0.69	-	0.88	-	-	1.0	-	-
282	Corn	Br	9/17	NA	3.2	-	-	-	-	-	-	-	-	-
283	"	Br	9/17	NA	2.9	-	-	-	-	-	-	-	-	-
284	"	-Replicate	9/17	NA	2.3	-	-	-	-	-	-	-	-	-
285	"	Samples	Br	9/17	NA	2.7	-	-	-	-	-	-	-	-
286	"	Samples	Br	9/17	NA	2.8	-	-	-	-	-	-	-	-
311	Milo	Hr	9/24	NA	3.1	1.6	0.12	-	-	-	-	2.1	-	-
Ringold Farms														
173	Wheat	Ke	7/22	NA	7.7	4.1	0.86	-	1.3	-	-	0.96	0.052	-
248	"	Me	9/9	NA	5.6	2.4	0.70	0.15	-	-	-	-	-	-
262	Wheat	Pr	9/17	NA	3.8	1.4	-	-	-	-	0.82	-	0.014	-
263	"	Pr	9/17	NA	3.4	1.2	-	-	-	-	0.97	-	-	-
264	"	- Replicate	Pr	9/17	NA	3.4	1.3	-	-	-	0.50	1.0	-	0.006
265	"	Samples	Pr	9/17	NA	3.9	1.4	-	-	-	-	1.3	-	-
266	"	Samples	Pr	9/17	NA	4.0	1.1	-	-	-	-	2.8	0.010	-
Grain Elevator, Prosser, Washington														
61	Wheat 1957 Storage		2/26	NA	4.0	-	-	-	0.071	-	-	-	-	-
78	"		4/16	NA	3.0	-	-	-	0.063	-	-	-	-	-
79	"		4/16	NA	3.5	-	-	-	0.072	-	-	-	-	-
												Lost	0.012	0.008

Results less than the reporting limit are indicated by a (-). NA - Not Analyzed

APPENDIX D-3 (contd.)

Code	Crop	Reporting Limits*	Location Code	Date	Sc ⁴⁶	K ⁴⁰	Zn ⁶⁵	Zr ⁹⁵ -Nb ⁹⁵	Cs ¹³⁷	Fu ¹⁰³⁺	Fu ¹⁰⁶	I ¹³¹	Cr ⁵¹	Ce ¹⁴¹⁺	Ce ¹⁴⁴	Sr ⁸⁹	Sr ⁹⁰
					0.1	0.3	0.1	0.1	0.05	0.5	0.1	0.5	0.5	0.01	0.006		
<u>Benton City and Yakima Valley</u>																	
158 Wheat	Bc	7/16	NA	3.3	-	0.44	-	-	-	-	-	0.57	0.021	0.015			
159 Rye	Bc	7/16	" Lost	3.7	-	0.44	-	-	-	-	-	0.78	0.024	0.028			
170 Wheat	Bc	7/22	-	4.8	-	-	0.070	-	-	-	-	-	0.023	0.016			
322 Wheat	Bc	9/25	-	-	-	-	-	-	-	-	-	-	-	-	Analysis not Complete		
<u>Finley District</u>																	
185 Barley	Sr	7/22	NA	4.8	-	0.22	-	-	-	-	-	-	"	"	"	"	"
222 Corn	Cr	8/19	"	2.7	-	-	-	-	-	-	-	-	-	-	-	-	-
289 Milo	Hs	9/18	-	3.7	-	0.14	-	-	-	-	-	-	0.80	1.0	-	-	-
290 "	Hs	9/18	-	4.1	-	0.14	0.054	-	-	-	-	-	0.55	"	Not Analyzed	"	"
291 "	Replicate Samples	Hs	9/18	-	4.1	-	-	0.081	-	-	-	-	0.66	"	"	"	"
292 "		Hs	9/18	-	4.5	-	0.14	0.064	-	-	-	-	0.70	"	"	"	"
293 "		Hs	9/18	-	4.8	-	0.12	0.053	-	-	-	-	0.50	0.83	Analysis not Complete		
315 Milo	Hs	9/24	-	6.3	-	0.14	0.076	-	-	-	-	-	-	-	-	-	-
<u>Walla Walla, Washington</u>																	
343 Wheat	Kr	10/ 6	-	3.9	-	0.71	0.42	2.8	-	2.1	3.5	"	"	"	"	"	"
<u>Spokane, Washington</u>																	
359 Wheat	Cr	10/ 6	-	3.7	-	0.98	0.25	2.5	-	-	4.4	"	"	"	"	"	"
<u>Pendleton, Oregon</u>																	
340 Barley	Hn	10/ 6	-	4.2	-	-	0.081	-	-	-	1.0	-	-	-	Not Analyzed		
341 "	Hn	10/ 6	-	4.0	-	-	0.052	-	-	-	-	-	-	-	Analysis not Complete		
342 "	Hn	10/ 6	-	4.4	-	-	-	-	-	-	-	-	-	-	Not Analyzed		

* Results less than the reporting limit are indicated by a (-). NA - Not Analyzed

APPENDIX D-4

**CONCENTRATIONS OF SEVERAL RADIONUCLEIDES IN VEGETABLES PURCHASED FROM GROWERS
AT SELECTED LOCATIONS - 1959 IN UNITS OF $10^{-6} \mu\text{C}/\text{g}$**

APPENDIX D-4 (contd.)

APPENDIX D-4 (contd.)

<u>Code</u>	<u>Crop</u>	<u>Location</u>	<u>Date</u>	<u>Sc⁴⁶</u>	<u>K⁴⁰</u>	<u>Zn⁶⁵</u>	<u>Zr⁹⁵-Nb⁹⁵</u>	<u>Cs¹³⁷</u>	<u>Ru¹⁰³⁺</u>	<u>Ru¹⁰⁶</u>	<u>Cr⁵¹</u>	<u>Ce¹⁴⁴</u>	<u>Sr⁸⁹</u>	<u>Sr⁹⁰</u>
<u>Reporting Limits</u>		<u>Code</u>		<u>0.1</u>	<u>0.3</u>	<u>0.1</u>	<u>0.1</u>	<u>0.05</u>	<u>0.5</u>	<u>0.1</u>	<u>0.5</u>	<u>0.5</u>	<u>0.01</u>	<u>0.00</u>
<u>Walla Walla, Washington</u>														
347 Romaine	Oe	10/ 6	-	4.2	-	-	-	-	-	-	-	-	-	-
348 Cabbages	Sr	10/ 6	-	2.6	-	-	-	-	-	-	-	-	-	-
349 Radishes	Mo	10/ 6	-	3.1	-	-	-	-	-	-	-	-	-	-
<u>Spokane, Washington</u>														
350 Carrots	Sn	10/ 6	-	3.2	-	-	-	-	-	-	-	-	-	-
351 Lettuce	Ki	10/ 6	-	2.1	-	-	-	-	-	-	-	-	-	-
352 Turnips	Ta	10/ 6	-	1.4	-	-	-	-	-	-	-	-	-	-
<u>Local Purchase - California Grown</u>														
67 Carrots	3/31	NA	2.8	-	-	-	-	-	-	-	-	-	-	-
68 Carrot tops	3/31	II	2.9	-	3.0	-	-	-	0.51	-	-	-	-	-
70 Carrots	4/ 9	II	2.7	-	-	-	-	-	-	-	-	-	-	-
75 "	4/16	II	3.2	-	-	-	-	-	-	-	-	-	-	-
76 Carrot tops	4/16	II	6.6	0.11	-	-	-	0.095	-	-	-	-	-	2.2
86 Carrots	4/22	II	2.2	-	-	-	-	-	-	-	-	-	-	-
87 Carrot tops	4/22	II	-	-	-	-	-	4.1	-	-	-	-	-	4.1
108 Carrots	5/25	II	2.5	-	-	-	-	-	-	0.60	-	-	-	0.12

Results less than the reporting limits are indicated by a (-). NA - Not Analyzed

Analysis not Complete

" Not Analyzed

" Analysis not Complete

" Not Analyzed

" Analysis not Complete

" Not Analyzed

" Analysis not Complete

APPENDIX D-5

CONCENTRATIONS OF SEVERAL RADIONUCLIDES IN FRUITS PURCHASED FROM GROWERS
AT SELECTED LOCATIONS - 1959. IN UNITS OF $10^{-6} \mu\text{c}/\text{g}$.

Code	Crop	Reporting Limits	Location Code	Date	Sc 46	K 40	Zn 65	Zr 95-Nb 95	Cs 137	Ru 103+ Ru 106	Ru 131	Cr 51	Ce 141+ Ce 144	Sr 89	Sr 90
<u>Riverview Irrigation District</u>															
122	Strawberries	Zn	6/29	NA	1.8	0.17	-	-	-	-	-	0.90	-	-	-
148	Boysenberries	Pr	7/16	"	3.0	0.23	-	-	-	-	-	-	-	-	-
167	Apricots	Hs	7/22	"	5.5	-	0.21	-	-	-	-	-	-	0.016	0.0067
168	Apples	Hs	7/22	"	1.4	-	-	-	-	-	-	-	-	-	-
217	Peaches	An	8/19	"	2.4	-	-	-	-	-	-	-	-	-	-
267	Grapes	Ar	9/17	-	2.5	-	-	-	-	-	-	-	-	-	-
268	"	- Replicate	Ar	9/17	-	2.4	-	-	-	-	-	-	-	-	-
269	"	Samples	Ar	9/17	-	2.4	-	-	-	-	-	-	-	-	-
270	"	Ar	9/17	-	2.3	-	-	-	-	-	-	-	-	-	-
312	Pears	Ar	9/25	-	2.6	-	-	-	-	-	-	-	-	-	-
<u>Ringold Farms</u>															
150	Peaches	KY	7/16	NA	2.3	-	-	-	-	-	-	-	-	-	-
151	Apricots	KY	7/16	"	2.8	-	-	-	-	-	-	-	-	-	-
171	Peaches	KY	7/22	"	2.0	0.11	-	-	-	-	-	-	-	-	-
172	Apricots	KY	7/22	"	2.7	-	-	-	-	-	-	-	-	-	0.016
195	"	KY	7/22	"	2.7	-	-	-	-	-	-	-	-	-	-
196	"	KY	7/22	"	5.3	-	-	-	-	-	-	-	-	-	-
212	Pears	KY	7/22	"	1.6	-	-	-	-	-	-	-	-	-	-
213	Apples	KY	8/19	"	1.5	-	-	-	-	-	-	-	-	-	-
214	Plums	KY	8/19	"	2.5	-	-	-	-	-	-	-	-	-	-
215	Peaches	KY	8/19	"	2.5	-	-	-	-	-	-	-	-	-	-
308	Grapes	Pr	9/24	-	2.7	-	-	-	-	-	-	-	-	-	-
<u>Benton City and Yakima Valley</u>															
156	Peaches	Mg	7/17	NA	3.8	-	-	-	-	-	-	-	-	-	0.012
157	Apricots	Dr	7/16	"	2.8	-	-	-	-	-	-	-	-	-	0.0081
177	"	Er	7/22	"	3.1	-	-	-	-	-	-	-	-	-	0.011
319	Apples	Se	9/25	-	1.8	-	-	-	-	-	-	-	-	-	0.0082
															Not Analyzed

Results less than the reporting limit are indicated by a (-). NA - Not Analyzed

APPENDIX D-5 (contd.)

Results less than the reporting limit are indicated by a (-).

APPENDIX D-6

CONCENTRATIONS OF SEVERAL RADIONUCLIDES IN ALFALFA, HAY AND PASTURE GRASS PURCHASED FROM GROWERS AT SELECTED LOCATIONS - 1959. IN UNITS OF $10^{-6} \mu\text{C}/\text{g}$.

Location Code	Crop Code	Reporting Limits	Date	Sc ₄₆	K ₄₀	Zn ₆₅	Zr ₉₅ -Nb ₉₅	Cs ₁₃₇	Ru ₁₀₃₊	Ru ₁₀₆	Cr ₅₁	Ce ₁₄₁₊	Ce ₁₄₄	Sr ₈₉	Sr ₉₀
Riverview Irrigation District															
80 Fresh Alfalfa	An	4/22	N/A	5.3	0.48	9.4	0.40	2.3	-	-	3.1	-	6.8	0.20	0.023
92 "	An	4/29	"	8.2	1.4	3.7	0.48	0.99	-	-	-	2.1	0.19	0.032	
93 "	An	5/6	"	10.	0.65	9.5	0.45	2.6	-	-	3.8	"	Not Analyzed	"	
99 "	An	5/13	"	3.4	0.24	1.9	0.13	0.50	-	-	-	6.7	-	-	
102 "	An	5/20	"	6.8	0.93	6.5	0.53	1.1	-	-	-	2.0	0.13	0.022	
106 "	Hs	5/25	"	6.0	0.18	2.7	0.21	-	-	-	2.8	0.72	-	0.072	
127 "	An	7/8	"	4.7	1.2	0.22	0.10	-	-	-	10	1.4	0.020	0.025	
142 "	An	7/15	"	3.0	3.1	2.0	-	1.1	-	-	-	"	Not Analyzed	"	
143 "	Hs	7/16	"	6.3	0.12	0.15	-	-	-	-	-	-	"	"	
189 "	Hs	8/10	"	6.3	-	-	-	-	-	-	-	-	"	"	
202 "	An	8/12	"	6.7	0.36	-	-	-	-	-	1.2	-	0.67	"	
211 "	Hs	8/19	"	5.7	0.10	0.13	-	-	-	-	-	"	"	"	
218 "	An	8/19	"	7.8	2.2	0.17	-	0.85	0.27	8.9	-	3.5	0.093	-	
231 "	An	8/26	"	9.7	2.2	0.24	0.057	0.52	-	7.6	2.7	2.7	Not Analyzed	"	
233 Pasture Grass	Pr	8/27	"	10	8.0	0.49	0.14	-	-	25	-	-	-	-	
240 Fresh Alfalfa	Pr	9/1	-	6.4	1.1	-	-	-	-	-	15	-	0.051	0.022	
242 "	An	9/2	0.16	5.6	2.8	-	-	-	-	-	10	-	-	0.036	
249 "	An	9/10	0.28	5.6	1.8	-	-	-	-	-	31	12	-	0.020	
251 "	An	9/16	-	5.8	0.90	0.15	-	0.85	1.0	-	-	-	-	Not Analyzed	
272 "	Pt	9/17	-	6.2	-	0.24	0.073	-	-	-	-	1.4	"	"	
273 "	- Replica Pt Samples		9/17	-	6.1	-	0.26	0.056	1.0	-	-	2.5	"	"	
274 "	Pt	9/17	0.16	6.4	-	0.18	-	0.81	-	-	-	3.0	"	"	
275 "	Pt	9/17	-	6.5	-	0.29	-	0.62	1.2	-	-	4.2	"	"	
276 "	Pt	9/17	-	6.6	-	0.22	0.080	0.76	-	-	-	2.4	"	"	
304 Alfalfa	An	9/23	0.34	11	0.36	0.36	0.067	1.7	-	-	32	1.3	Lost	Lost	
310 Fresh Alfalfa	Pt	9/24	-	5.8	1.0	0.40	0.22	1.4	-	-	1.6	3.2	Lost	Lost	
328 "	An	9/30	0.23	8.1	2.2	0.20	-	0.85	1.3	3.7	1.8	1.3	Lost	Not Analyzed	
363 "	An	10/4	0.27	6.0	2.2	0.20	0.10	1.1	-	1.3	3.7	Analysis not Complete	"	"	
364 "	An	10/21	0.18	6.3	1.7	0.26	0.061	0.28	-	7.5	1.9	1.9	Lost	Analysis not Complete	
365 "	Pt	10/23	-	4.1	-	0.43	0.20	1.4	-	2.5	0.22	-	Lost	"	
374 "	An	10/28	-	5.3	0.17	0.18	0.060	1.7	-	-	-	1.7	1.9	Analysis not Complete	"

Results less than the reporting limit are indicated by a (-). NA - Not Analyzed

APPENDIX D-6 (contd.)

<u>Code</u>	<u>Crop</u>	<u>Location Code</u>	<u>Date</u>	<u>Sc 46</u>	<u>K 40</u>	<u>Zn 65</u>	<u>Zr 95-Nb 95</u>	<u>Cs 137</u>	<u>Ru 103+</u>	<u>Ru 106</u>	<u>I-131</u>	<u>Cr 51</u>	<u>Ce 141+</u>	<u>Ce 144</u>	<u>Sr 89</u>	<u>Sr 90</u>
	<u>Reporting Limits</u>			0.1	0.3	0.1	0.1	0.05	0.5	0.1	0.5	0.5	0.5	0.5	0.01	0.006
	<u>Ringold Farms</u>															
216	Fresh Alfalfa	Pr	8/19	NA	7.2	0.16	0.13	-	-	-	0.39	-	0.89	-		
257	"	Pr	9/17	0.15	6.5	1.4	0.17	-	-	-	5.0	-	0.073	-		
258	"	Replicate	9/17	-	5.8	1.3	0.15	-	-	0.37	3.5	0.77	0.052	0.008		
259	"	Samples	9/17	0.13	5.6	1.4	0.15	-	-	-	4.4	1.3	0.50	0.017		
260	"	Pr	9/17	0.20	6.4	1.7	0.13	-	-	-	6.0	0.70	0.074	-		
261	"	Pr	9/17	0.13	6.4	1.2	0.15	0.079	0.67	0.58	2.3	4.2	Analysis not Complete			
316	Fresh Alfalfa	Pr	9/24	0.14	12	2.1	0.21	0.054	1.2	-	7.2	1.7	Lost	Lost		
367	Fresh Alfalfa	Ke	10/23	0.16	5.3	0.32	0.20	0.20	1.1	0.27	1.8	1.2	Analysis not Complete			
	<u>Benton City and Yakima Valley</u>															
104	Fresh Alfalfa	Gv	5/22	NA	5.8	0.11	2.5	0.28	0.84	-	-	-	1.4	-		
105	"	Ss	5/22	"	7.4	0.15	3.7	0.27	0.70	-	-	-	2.7	Not Analyzed		
153	Dry Hay	Bc	7/22	"	24	-	3.8	-	2.7	-	-	-	6.4	0.14	0.19	
179	"	Bc	7/22	"	23	-	1.7	-	-	-	-	-	2.3	-	Lost	
321	Fresh Alfalfa	Bc	9/25	-	9.6	-	0.12	-	0.53	-	-	-	0.87	Analysis not Complete		
368	"	Bc	10/23	-	5.2	-	0.14	0.14	0.82	0.27	0.98	1.1	Lost	Lost		
378	"	Bc	11/2	0.18	4.8	-	0.17	0.17	0.96	0.32	1.0	1.7	Analysis not Complete			
	<u>Finley District</u>															
103	Fresh Alfalfa	Sp	5/22	NA	6.0	-	6.0	0.43	1.0	-	-	-	4.9	-	0.35	
140	"	Sp	7/10	"	4.2	-	3.32	-	-	-	-	-	0.62	0.037	0.071	
162	Dry Hay	Sp	7/17	"	21	-	3.4	-	2.5	-	-	-	2.5	0.290		
187	Fresh Alfalfa	Sp	7/22	"	8.5	-	-	-	-	-	-	-	-	0.019		
225	"	Sp	8/19	"	7.3	-	0.27	-	-	-	-	-	-	0.036	0.021	
294	Fresh Alfalfa	Sp	9/19	-	5.5	-	0.59	0.12	1.1	-	-	-	1.9	0.037	0.040	
295	"	Replicate	9/18	-	12	-	0.40	0.45	1.5	-	-	-	2.0	Not Analyzed	"	
296	"	Samples	9/18	-	14	-	0.44	0.17	1.7	-	-	-	3.7	"	"	
297	"	Sp	9/18	-	17	-	-	-	-	-	-	-	3.0	"	"	
298	"	Sp	9/18	-	14	-	0.46	0.15	1.8	-	-	-	3.4	"	"	
313	"	Sp	9/24	-	10	-	1.2	0.21	2.4	-	-	-	5.9	"	"	
366	Fresh Alfalfa	Sp	10/23	-	4.9	-	0.30	0.14	1.5	0.15	-	-	2.1			

Results less than the reporting limit are indicated by a (-). NA - Not Analyzed

APPENDIX D-6 (contd.)

Results less than the reporting limit are indicated by a (-).

APPENDIX D-7

CONCENTRATIONS OF SEVERAL RADIONUCLIDES
IN MISCELLANEOUS LOCALLY PURCHASED FOODSTUFFS* - 1959

Code	Type	Location Code	Date	Sc ⁴⁶	K ⁴⁰	Zn ⁶⁵	Zr ⁹⁵ -Nb ⁹⁵	Cs ¹³⁷	Ru ¹⁰³⁺	Ru ¹⁰⁶	I ¹³¹	Cr ⁵¹	Ce ¹⁴¹⁺ Ce ¹⁴⁴	
				0.1	0.3	0.1	0.1	0.05	0.5	0.1	0.5	0.5	0.5	
<u>Reporting Limits</u>														
Meat														
388	Ground Beef	Brand A	11/23	-**	2.0	-	0.32	-	-	-	-	-	-	-
390	Ground Beef	Brand B	11/23	-	1.8	-	0.32	-	-	-	-	-	-	-
394	Ground Beef	Brand C	11/23	-	2.2	-	-	0.064	-	-	-	-	-	-
396	Ground Beef	Brand D	11/23	-	2.6	-	0.33	-	-	-	-	-	-	-
<u>Canned Baby Food - Strained</u>														
429	Pork	Brand A	12/10	-	1.8	-	-	-	0.082	-	-	-	-	-
430	Beef	Brand A	12/10	-	2.0	-	-	-	0.072	-	-	-	-	-
431	Lamb	Brand A	12/10	-	1.6	-	-	-	0.13	-	-	-	-	-
432	Chicken	Brand A	12/10	-	0.82	-	-	-	-	-	-	-	-	-
434	Peas	Brand A	12/10	-	0.67	-	-	-	-	0.61	-	-	-	-

* Strontium analyses are not complete for meat and canned baby food.

** Results less than reporting limit are indicated by a (-).

INTERNAL DISTRIBUTION

Copy Number

1	G. E. Backman
2	G. D. Brown
3	B. E. Clark, Jr.
4	J. J. Davis
5	R. F. Foster
6	W. C. Hanson
7	J. W. Healy
8	F. E. Holt
9	P. C. Jerman
10	R. L. Junkins
11	A. R. Keene
12	M. W. McConiga
13	I. C. Nelson
14	J. M. Nielson
15	W. C. Roesch
16	A. J. Stevens
17	C. M. Unruh
18	J. W. Vanderbeek
19	E. C. Watson
20	300 Files
21	Record Center
22 - 25	G. E. Technical Data Center, Schenectady
26 - 157	Extra

EXTERNAL DISTRIBUTION

Copy Number

3	Aberdeen Proving Ground
1	Aerojet-General Corporation
1	Aerojet-General, San Ramon (I00-880)
1	AFPR, Boeing, Seattle
3	AFPR, Lockheed, Marietta
2	Air Force Special Weapons Center
2	ANP Project Office, Convair, Fort Worth
1	Alco Products, Inc.
1	Allis-Chalmers Manufacturing Company
3	Argonne Cancer Research Hospital
10	Argonne National Laboratory
1	Armed Forces Special Weapons Project, Washington
4	Army Chemical Center
1	Army Chemical Center (Taras)
1	Army Chemical Corps
1	Army Medical Research Laboratory
1	Army Signal Research and Development Laboratory
1	Atomic Bomb Casualty Commission
1	AEC Scientific Representative, Japan
3	Atomic Energy Commission, Washington
3	Atomics International
2	Babcock and Wilcox Company (NY00-1940)
2	Battelle Memorial Institute
2	Bettis Plant
4	Brookhaven National Laboratory
2	Brooks Army Medical Center
1	Brush Beryllium Company
1	BAR, Goodyear Aircraft, Akron
1	BAR, Grumman Aircraft, Bethpage
1	Bureau of Medicine and Surgery
1	Bureau of Mines, Albany
1	Bureau of Mines, Salt Lake City
1	Bureau of Ships (Code 1500)
1	Bureau of Yards and Docks
2	Chicago Operations Office
1	Chicago Patent Group
1	Columbia University (Failla)
1	Combustion Engineering, Inc.
1	Committee on the Effects of Atomic Radiation
1	Convair-General Dynamics Corporation, San Diego
3	Defense Research Member
2	Department of the Army, G-2

EXTERNAL DISTRIBUTION (contd.)

Copy Number

1	Division of Raw Materials, Washington
1	Dow Chemical Company (Rocky Flats)
3	duPont Company, Aiken
1	duPont Company, Wilmington
1	Edgerton, Germeshausen and Grier, Inc., Boston
1	Edgerton, Germeshausen and Grier, Inc., Las Vegas
1	Frankford Arsenal
2	General Electric Company (ANPD)
2	General Electric Company, St. Petersburg
1	Gibbs and Cox, Inc.
1	Goodyear Atomic Corporation
1	Grand Junction Operations Office
1	Hawaii Marine Laboratory
1	Iowa State College
3	Knolls Atomic Power Laboratory
2	Los Alamos Scientific Laboratory
1	Lovelace Foundation
1	M & C Nuclear, Inc.
1	Mallinckrodt Chemical Works
1	Maritime Administration
1	Martin Company
1	Massachusetts Institute of Technology (Hardy)
1	Mound Laboratory
1	National Academy of Sciences
1	National Aeronautics and Space Administration, Cleveland
2	National Bureau of Standards
1	National Cancer Institute
1	National Industrial Conference Board
1	National Lead Company, Inc., Winchester
1	National Lead Company of Ohio
1	National Library of Medicine
1	Naval Medical Research Institute
3	Naval Research Laboratory
1	New Brunswick Area Office
2	New York Operations Office
1	Nuclear Development Corporation of America
1	Oak Ridge Institute of Nuclear Studies
15	Office of Naval Research
1	Office of Naval Research (Code 422)
1	Office of Ordnance Reserach
1	Office of the Chief of Naval Operations
1	Office of the Surgeon General
1	Olin Mathieson Chemical Corporation
1	Ordnance Tank-Automotive Command

EXTERNAL DISTRIBUTION (contd.)

Copy Number

1	Patent Branch, Washington
6	Phillips Petroleum Company (NRTS)
1	Power Reactor Development Company
3	Pratt and Whitney Aircraft Division
1	Princeton University (White)
2	Public Health Service
1	Public Health Service, Savannah
1	Rensselaer Polytechnic Institute
1	Sandia Corporation, Albuquerque
1	Schenectady Naval Reactors Operations Office
1	Sylvania Electric Products, Inc.
1	Technical Research Group
1	Tennessee Valley Authority
3	The Surgeon General
2	Union Carbide Nuclear Company (ORGDP)
5	Union Carbide Nuclear Company (ORNL)
1	Union Carbide Nuclear Company (Paducah Plant)
1	USAF Project RAND
1	U. S. Geological Survey, Naval Gun Factory
1	U. S. Naval Ordnance Laboratory
1	U. S. Naval Postgraduate School
2	U. S. Naval Radiological Defense Laboratory
1	University of California at Los Angeles
3	University of California, Berkeley
2	University of California, Livermore
1	University of California, San Francisco
1	University of Chicago, USAF Radiation Laboratory
1	University of Puerto Rico
1	University of Rochester
1	University of Tennessee
1	University of Utah
1	University of Washington (Donaldson)
1	Walter Reed Army Medical Center
1	Watertown Arsenal
1	Western Reserve University
1	Westinghouse Electric Corporation (Schafer)
6	Wright Air Development Center
1	Yankee Atomic Electric Company
325	Technical Information Service Extension
100	Office of Technical Services, Washington

EXTERNAL DISTRIBUTION (contd.)

Copy Number

1	L. B. Dworsky - Public Health Service, Portland, Oregon
1	C. M. Everts - Oregon State Board of Health, Portland
1	A. Garton - Washington Pollution Control Commission, Olympia, Washington
1	E. C. Jensen - Washington State Dept. of Health, Seattle, Washington
1	E. F. Miller - AEC - WASH. Division of Production
1	K. L. Englund - AEC-HOO
1	C. F. Whetsler - City Water Superintendent of Pasco 412 W. Clark, Pasco, Washington
1	A. W. Klement, Jr. - Fallout Studies Branch Division of Biology and Medicine, AEC, Washington 25, D. C.
1	Dr. T. S. Ely - Office of Health and Safety, AEC, Washington 25, D. C.
1	A. T. Neale - Pollution Control Commission, Olympia, Washington